arXiv:1101.0424v1 [math.NT] 2 Jan 2011

Two-dimensional Idèles with Cycle Module Coefficients

O. Braunling

Abstract. We give a flasque resolution of the Zariski sheaves of Rost cycle modules in terms of 2-dimensional idèles on surfaces. We give explicit formulas relating this idèle resolution with the Gersten-type and Čech resolution. As an application we recover an old formula expressing intersection multiplicities in terms of tame symbols and a Čech analogue thereof. We show that this approach is workable for explicit computations.

Leaving details and definitions to the main body of this text, our main result can be summarized by saying that for integral smooth surfaces over a field k and a Rost cycle module M_* there is a flasque resolution of Zariski sheaves

$$0 \to \mathcal{M}_* \to \mathbf{I}_0 \to \mathbf{I}_1 \to \mathbf{I}_2 \to 0, \tag{0.1}$$

where $\mathcal{M}_*$ is the Zariski sheaf associated to M_* and $\mathbf{I}_\bullet$ are sheaves of 'surface idèles' - a large part of the text works out what this is supposed to mean. As a consequence one obtains subquotients of 'idèle objects' which express for example

- K-cohomology groups $H^p(X, \mathcal{K}_q)$, the terms on the E_2-page of the Brown-Gersten-Quillen coniveau spectral sequence;
- in particular Chow groups $H^p(X, \mathcal{K}_p) \cong \mathrm{CH}^p(X)$;
- cohomology groups $H^p_{\mathrm{Zar}}(X, \mathcal{H}^q(\mu_n^{\otimes r}))$, $(n, \operatorname{char} k) = 1$, the terms of the E_2-page of the Bloch-Ogus coniveau spectral sequence ("$\mathcal{H}$-cohomology");
- certain variations of these, e.g. a variant $H^p(X, \mathcal{K}^t_q)$ of K-cohomology coming from Milnor K-groups modulo divisible elements; this is relevant if one wants to use so-called topological Milnor K-groups K_n^{top} in the idèles, see §2.3 for details.

These variations all stem from the basic principle that all these groups are sheaf cohomology groups of Zariski sheaves of various Rost cycle modules.

The direct 1-dimensional analogue of this theory with K-theory coefficients in degree 1 would recover either Chevalley's classical idèles (without infinite places) in the number field case or the corresponding function field analogue due to Iwasawa. Our idèle resolution, eq. 0.1, respects product structures[1] and as an application we recover a formula of expressing intersection multiplicities through tame symbols (and analogous explicit formulas for all cycle modules with a product structure). We give an example computation (a negative self-intersection number, see §8).

1991 *Mathematics Subject Classification.* 14F43, 19F27, 11R56, 14C17, 19E15.

University of Nottingham; supported by the EPSRC.

[1] a pairing of a cycle module with itself to itself

0.1. Applications. A bit stronger even, we formulate two bicomplexes whose E^1-pages (depending on the filtration chosen) recover either the Gersten-type resolution [**26**, proof of Cor. 6.5], whose cohomology is denoted by $A^p(X, M_*)$; the Čech resolution, whose cohomology is denoted by $\check{H}^p(X, \mathcal{M}_*)$; or the idèle resolution of $\mathcal{M}_*$ introduced in this text, whose cohomology will be denoted by $H^p(\Gamma(X, \mathbf{I}^\bullet))$ - the latter is suppressing its dependency on M_* for the sake of a convenient notation. We use this to determine *explicit* formulas for the comparison maps between explicit cocycle representatives in the respective resolutions:

$$\begin{tikzcd}
 & \text{Gersten} \arrow[dl] \arrow[dr] & \\
\text{Čech} \arrow[rr] & & \text{Idèles}
\end{tikzcd}$$

As an application of these comparison maps we obtain explicit formulas for the intersection pairing (à la Bloch, [**4**, Rmk. 2.14])

$$\mathrm{CH}^1(X) \otimes \mathrm{CH}^1(X) = H^1(X, \mathcal{K}_1) \otimes H^1(X, \mathcal{K}_1) \longrightarrow H^2(X, \mathcal{K}_2) = \mathrm{CH}^2(X)$$

in the Čech and idèle resolution (for Gersten this is well-known and gives the classical definition of the product of the Chow ring). This recovers a 30-year old formula of Parshin which expresses the intersection multiplicity in terms of tame symbols of idèles, cf. [**24**] (see Gorchinskiy [**12**] for another approach). We complement this by a further variation of this theme by using the comparison to the Čech resolution, giving

$$D \cdot D' = \sum_{x,y} j_* \partial_x^y \partial_y^\eta \{ f_{\beta_\eta \beta_y}, f'_{\beta_y \beta_x} \} \quad \text{(finite sum over tame symbols)} \tag{0.2}$$

(notation explained in the text). Here the indices β_* depend on a disjoint decomposition of the scheme space X related to the construction of an explicit contracting homotopy for the Čech complex. This (and the idèle) formula works for *any* intersection without needing the divisors to intersect properly (no "moving").

Leitfaden:

- We define 2-dimensional idèles with cycle module coefficients, Def. 3 on page 21;
- We prove this provides a flasque resolution of the Zariski sheaf $\mathcal{M}_*$ of a cycle module, Prop. 5; we denote these 'idèle representatives' of cohomology groups by $H^p(\Gamma(X, \mathbf{I}_\bullet))$;
- We give a bicomplex whose E^1-pages depending on the filtration chosen give either the Gersten resolution or our idèle resolution of $\mathcal{M}_*$, Prop. 6;
- As an application of that we obtain explicit formulas for mapping explicit cohomology class representatives from one resolution to another, specifically $A^1(X, M_*) \to H^1(\Gamma(X, \mathbf{I}_\bullet))$, Prop. 7, and $H^2(\Gamma(X, \mathbf{I}_\bullet)) \to A^2(X, M_*)$, Prop. 8;
- Using the classical Čech resolution for sheaf cohomology, there are similar bicomplexes whose E^1-pages depending on the filtration chosen give either the Gersten, idèle or Čech resolution. As an application of that we obtain quite analogous explicit formulas $\check{H}^1(X, \mathcal{M}_*) \to H^1(\Gamma(X, \mathbf{I}_\bullet))$, Prop. 9, and $\check{H}^2(X, \mathcal{M}_*) \to A^2(X, M_*)$, Prop. 10;
- More such formulas could be developed, but these are the only ones we have further use of;

- Next, we assume the cycle module M_* is equipped with a product structure $M_* \times M_* \to M_*$. We show that the cup product '$\smile$' in Čech cohomology has a counterpart in the idèle resolution, which we denote by '$\vee$' to have a clear distinction, an 'idèle cup product'. We prove that both induce the same product to cohomology, Prop. 11;
- Connecting this with the fact that the intersection product can be expressed as such a product in cohomology, we obtain an idèle and a Čech resolution formula for intersection multiplicities in terms of tame symbols. While the idèle version is known since Parshin, the latter seems to be new.

0.2. Acknowledgements. I would like to thank P. Arndt, J. P. Demailly, B. Kahn and A. Vishik for answering some questions I had. Moreover, K. Ardakov for spotting a crucial mistake which then led to various simplifications. F. Trihan, M. Morrow and I. Fesenko for their careful reading of earlier versions of this text and very, very useful conversations. Finally, more thanks to I. Fesenko for letting me work on this interesting question and providing the ideas of the bigger picture/perspective from which one should look at higher idèles (and beyond).

1. Overview of Multidimensional Adèles

This section surveys the ideas and motivations behind the concept of multidimensional adèles/idèles. It puts this work into a context, but is not necessarily required in any of the ensuing sections.

(1) *Origins:* The concept of (ordinary 1-dimensional) adèles/idèles has proven very successful for studying global fields, most notably number fields. They provide the natural backdrop for the formulation of global class field theory à la Chevalley, their topology reflects the basic pillars of algebraic number theory (e.g. finiteness of the class number, Dirichlet's Unit Theorem), and exploiting the self-duality of the adèles as a LCA group and the associated harmonic analysis one obtains a very powerful tool in the study of zeta and L-functions via zeta integrals on the adèles (known as the Tate-Iwasawa method, "Tate's Thesis", [**28**]). Especially the latter had previously only been available through works of Hecke on theta functions; a complicated immanently global approach not allowing for a global-local perspective as the adèle method does.

(2) *Idèles as Sheaves:* Let us focus on the function field case for a start. Then from a scheme perspective, adèles/idèles can be formulated as Zariski sheaves on an integral smooth curve X of finite type over a (finite[2]) field k. Let K denote its rational function field. Then adèles/idèles provide flasque resolutions of the sheaves $\mathcal{O}$ (adèles) and $\mathcal{O}^\times$ (idèles). For example, the classical formula recovering the ideal class group $\operatorname{Pic} \mathcal{O}_K$ from the idèle class group $\mathbf{C}_K$,

$$\operatorname{Pic} X = H^1(X, \mathcal{O}_X^\times) = \mathbf{C}_K / \prod \widehat{\mathcal{O}}_v^\times, \tag{1.1}$$

is just an immediate consequence of the flasque resolution

$$0 \longrightarrow \mathcal{O}^\times \xrightarrow{\text{diag}} \underline{K}^\times \oplus \prod \widehat{\mathcal{O}}_v^\times \xrightarrow{\text{diff}} \prod{}' \widehat{K}_v^\times \longrightarrow 0, \tag{1.2}$$

---

[2] Our discussion is valid for arbitrary residue fields. However, in general in order to have the whole adèle theory for curves available one needs a finite residue field for otherwise the adèles are not locally compact and then conventional harmonic analysis in not available. This is spiritedly discussed by A. Weil in [**31**, p. VI (Foreword), 2nd paragraph].

where $\underline{K}^{\times}$ is the locally constant sheaf with value $K^{\times}$ and the products are to be read as sheaves $U \mapsto \prod_v(\ldots)$ over all closed points $v \in U_{(0)}$ and the prime ′′ in $\prod'$ denotes a certain finiteness constraint. A nice exposition using this sheaf-theoretic picture is given in [**2**, §3], which in turn is based on Tate's work [**29**]. The latter note in particular gives an extremely elegant and short proof of the residue theorem $\sum_{x\in X_{(0)}} \operatorname{res}_x \omega = 0$ for a global rational 1-form $\omega \in \Omega^1_{K/\mathbf{F}_q}$ based on a genuinely adelic approach.

(3) *Adèles of Schemes:* Replacing multiplicative groups by additive groups, the counterpart of eq. 1.2 yields a quotient encoding $H^1(X, \mathcal{O}_X)$. Parshin then more generally saw the possibility to use variations of adèles to resolve arbitrary quasi-coherent sheaves for curves, and then also for surfaces [**23**]. Initially, these works were inspired by applications to class field theory and for example the ideas in the book [**27**] by Serre. There is a recent approach to a reciprocity law of Parshin for surfaces using 2-dimensional adèles by Osipov and Zhu [**21**]. The generalization of adèles to dimension 2 requires various new ideas. See [**3**], [**15**] for Beilinson's generalization allowing to resolve quasi-coherent sheaves using higher-dimensional adèles in any dimension. Generally, these idèles have a 'semi-simplicial' organization, as explained nicely in [**10**], [**11**], [**24**]; this will also be the case for the idèles in this text.

(4) *Coefficient Systems:* In order to generalize the multiplicative flasque resolution of eq. 1.2 a first observation is that for local domains $\mathcal{O}$ one has $K_1(\mathcal{O}) = \mathcal{O}^{\times}$ and if one represents elements of $K_2(\mathcal{O})$ as symbols $\sum\{a_i, b_i\}$ with $a_i, b_i \in \mathcal{O}^{\times}$ (this is possible by a theorem of v. d. Kallen if the residue field of $\mathcal{O}$ has ≥ 6 elements, [**30**], let us assume we are in this situation), one can write down the sequence eq. 1.2 with K_2 in place of K_1 and take the maps as in eq. 1.2, but this time applied to the two slots a_i, b_i individually. It is natural to ask whether the sequence remains exact. It turns out it does, i.e.

$$(1.3) \qquad 0 \longrightarrow \mathcal{K}_2 \longrightarrow \underline{K_2(K)} \oplus \prod\nolimits_{x\in U^{(1)}} K_2(\widehat{\mathcal{O}}_x) \xrightarrow{d'} \prod\nolimits'_{x\in U^{(1)}} K_2(\widehat{K}_x) \longrightarrow 0$$

with $d' : (x, y_x) \mapsto y_x - \operatorname{res}^{\widehat{K}_x}_K x$, $\widehat{K}_x := \operatorname{Frac} \widehat{\mathcal{O}}_x$ is a flasque resolution. For example, for the projective line $\mathbf{P}^1_k$ this would imply

$$(1.4) \qquad H^1(\mathbf{P}^1_k, \mathcal{K}_2) = \frac{\prod'_{x\in(\mathbf{P}^1_k)^{(1)}} K_2(\widehat{K}_x)}{d'(K_2(K) \oplus \prod_{x\in(\mathbf{P}^1_k)^{(1)}} K_2(\widehat{\mathcal{O}}_x))} \qquad (\cong k^{\times}).$$

We will not prove this (not too difficult fact) in this text as we wish to focus on surfaces. The group $H^1(\mathbf{P}^1_k, \mathcal{K}_2)$ is an example of a K-cohomology group, in sheaf theory formulation. Generally, the K-cohomology groups $H^p(X, \mathcal{K}_q)$ constitute the E_2-page of the Brown-Gersten-Quillen spectral sequence converging to the algebraic K-theory of a variety X (filtered by coniveau).

This leads to the question whether it is generally possible to resolve the K-theory sheaves $\mathcal{K}_q$ in the style of eq. 1.2. If possible, this then naturally leads to many more subquotient formulas for K-cohomology groups in the spirit of eq. 1.4. We affirmatively answer this in the present work for smooth surfaces as a special case of our results (because the algebraic K-theory sheaves $\mathcal{K}_q$ are Zariski sheaves coming from a cycle module, see §2).

See also the work of Gorchinskiy [**12**] for many positive results in this direction for resolving certain homology sheaves (e.g. also cycle module sheaves, K-theory) in any dimension from a different perspective, however using adèles which replace completions by localizations (e.g. $\mathbf{Z}_{(p)}$ instead of the p-adics $\mathbf{Z}_p$), so-called "non-complete/rational adèles". See also the work of Osipov [**20**] for studies of the functorial behaviour of a certain notion of (completed) K-theory idèles. These works were very inspiring.

2. Cycle Modules

All schemes are assumed Noetherian, separated and finite-dimensional. Fix a field k.

Firstly, let us recall the basic notions of Rost's theory of cycle modules, see [**26**] for details (or [**7**] for some newer developments). We summarize the key features, mildly less general as in Rost's setup.

Rost works only with fields which are finitely generated over k as explained in [**26**, p328]. This is too restrictive for our purposes. We allow *all* field extensions F/k. Also, unlike in Rost's work, for us:

Definition 1. *A* (discrete) valuation *is a* $\mathbf{Z}$*-valued valuation, trivial on* k*.*

We should point out that with this definition of a (discrete) valuation even $k(s,t)$ possesses $\mathbf{Z}$-valued valuations with residue field k, i.e. the transcendence degree over k can drop further than just by 1. The corresponding valuation rings are Noetherian, but not essentially of finite type over k. Rost rules them out by demanding valuations to be geometric over k, but we cannot do this since this forbids most interesting valuations arising after completion, e.g. the t-adic valuation of $k((t))$ is not geometric over k.

Moreover, the p-adic valuations on $\mathbf{Q}$ are excluded by the above definition since they are not trivial on $\mathbf{Q}$.

2.1. Axiomatic Patterns.

For every field extension F/k let $K_*^M(F)$ denote the $\mathbf{Z}_{\geq 0}$-graded anticommutative Milnor K-theory ring over F,

$$K_*^M(F) = \coprod_{n\geq 0} K_n^M(F) \qquad\qquad = \mathbf{Z} \oplus F^\times \oplus K_2^M(F) \oplus \ldots .$$

Then a *cycle module* is an object function

$$\{\text{field extensions } F/k\} \longrightarrow \{\mathbf{Z}\text{-graded abelian groups}\}$$

along with some extra structure **D1-D4** satisfying various axioms. The extra structure (called *transfers*) is given as follows: Suppose E, F are fields containing k:

D1: For every field extension $(k \hookrightarrow)\ F \hookrightarrow E$ there is an abelian group morphism of relative degree 0,

$$\mathrm{res}_F^E : M_*(F) \longrightarrow M_*(E),$$

called *restriction*.

D2: For every *finite* field extension $(k \hookrightarrow)\ F \overset{\text{finite}}{\hookrightarrow} E$ there is an abelian group morphism of relative degree 0,

$$\mathrm{cor}_F^E : M_*(E) \longrightarrow M_*(F),$$

called *corestriction* (or *norm*).

D3: For every field ($k \hookrightarrow$) F the group $M_*(F)$ has the structure of a graded left-$K^M_*(F)$-module,

$$K^M_n(F) \cdot M_m(F) \subseteq M_{n+m}(F).$$

D4: For every field ($k \hookrightarrow$) F and every discrete valuation v on F, there is an abelian group homomorphism of relative degree -1,

$$\partial^F_v : M_*(F) \longrightarrow M_{*-1}(\kappa(v)),$$

where $\kappa(v)$ denotes the residue field of the valuation ring $A_v \subseteq F$ corresponding to v. ∂^F_v is called the *boundary map* at v.

Various axioms need to be satisfied, see below. For later use one also needs an elaboration of **D4**, which will play a very prominent role in this text:

Let $X \to \operatorname{Spec} k$ be an excellent scheme, $x \in X^{(i)}$ a codimension i point for some i, $y \in X^{(i+1)}$ a codimension $i+1$ point, $y \in \overline{\{x\}}$, i.e. y can also be read as a codimension 1 point in the integral closed subscheme $\overline{\{x\}} \hookrightarrow X$. One defines a *boundary map*

$$\partial^x_y : M_*(\kappa(x)) \longrightarrow M_{*-1}(\kappa(y))$$

as follows: The stalk $\mathcal{O}_{\overline{\{x\}},y}$ is 1-dimensional local ring, and a domain since $\overline{\{x\}}$ is integral. By excellence, the normalization

$$\varphi : \operatorname{Spec} \mathcal{O}'_{\overline{\{x\}},y} \longrightarrow \operatorname{Spec} \mathcal{O}_{\overline{\{x\}},y} \tag{2.1}$$

(in its own field of fractions $\kappa(x)$) is a finite morphism, so the maximal ideal y in $\operatorname{Spec} \mathcal{O}_{\overline{\{x\}},y}$ has a finite preimage $y'_1, \ldots, y'_r \in \operatorname{Spec} \mathcal{O}'_{\overline{\{x\}},y}$, precisely the maximal ideals of this semilocal normal 1-dimensional scheme. By normality the localizations $(\mathcal{O}'_{\overline{\{x\}},y})_{y'_i}$ are DVRs. Let $v_1, \ldots, v_r$ denote the corresponding discrete valuations of the fraction field $\kappa(x)$. Now one defines

$$\partial^x_y := \sum_{i=1}^{r} \operatorname{cor}^{\kappa(v_i)}_{\kappa(y)} \circ \partial^{\kappa(x)}_{v_i} : M_*(\kappa(x)) \longrightarrow M_{*-1}(\kappa(y)), \tag{2.2}$$

where $\kappa(v_i)$ denotes the residue field with respect to the valuation v_i (i.e. the residue field $\kappa(y'_i)$ of the semilocal ring $\mathcal{O}'_{\overline{\{x\}},y}$), $\partial^{\kappa(x)}_{v_i}$ is as in **D4**. Note that the finiteness of the normalization morphism φ, eq. 2.1, implies that $\kappa(v_i)/\kappa(y)$ is a *finite* field extension, so **D2** is indeed available.

These transfer morphisms need to satisfy a long list of axioms, **R1a-R3f** ("cycle premodule axioms"), for which we refer to [**26**, Def. 1.1], and the two crucial axioms ("cycle module axioms"):

FD: For every normal integral scheme $X \to \operatorname{Spec} k$ with rational function field K, and any $\alpha \in M_*(K)$, we have

$$\partial^K_x \alpha \neq 0 \in M_{*-1}(\kappa(x))$$

for at most finitely many codimension 1 points $x \in X^{(1)}$ (note that the stalk $\mathcal{O}_{X,x}$ at a codimension 1 point is a DVR with field of fractions K and residue field $\kappa(x)$, thus giving a discrete valuation v on K, so ∂^K_x refers to this valuation; equivalently use ∂^η_x, where η denotes the generic point of X).

C: For every 2-dimensional local domain $(R, \mathfrak{m})$ with field of fractions K, $\operatorname{Spec} R \to \operatorname{Spec} k$, and every $\alpha \in M_*(K)$ we have

$$\sum_x \partial_{\mathfrak{m}}^x \partial_x^{(0)} \alpha = 0 \in M_{*-2}(\kappa(\mathfrak{m})), \tag{2.3}$$

where x runs through all height 1 primes, i.e. $x \in (\operatorname{Spec} R)^{(1)}$, (0) denotes the generic point of $\operatorname{Spec} R$.

2.2. Unramified Cycles and Rost Complex. We need several other results and constructions related to cycle modules. Firstly, for every excellent n-equidimensional scheme $X \to \operatorname{Spec} k$ we define

$$C^p(X, M_q) := \coprod_{x \in X^{(p)}} M_{q-p}(\kappa(x)), \qquad \text{(also denoted by } C^p(X; M, q)) \tag{2.4}$$

where the coproduct is taken over the codimension p points of X. Here q is a $\mathbf{Z}_{\geq 0}$-grading and along $p \geq 0$ one obtains a complex $C^\bullet(X, M_q)$ by defining the q-degree-preserving differential

$$\partial_X : C^p(X, M_*) \longrightarrow C^{p+1}(X, M_*)$$

by using for all pairs $x \in X^{(p)}$ and $y \in X^{(p+1)}$, $y \in \overline{\{x\}}$ (i.e. y is a codimension 1 point of the integral closed subscheme $\overline{\{x\}}$), the boundary map

$$\partial_y^x : M_*(\kappa(x)) \longrightarrow M_{*-1}(\kappa(y)) \tag{2.5}$$

of eq. 2.2 and defining $\partial := \sum \partial_y^x$ for all such pairs $y \in \overline{\{x\}}$. This is a differential, $\partial_X^2 = 0$, see [**26**, Lemma 3.3]. In fact this property is essentially equivalent to axiom **C** of a cycle module. One defines

$$M_*(X) := \ker\left(\partial_X : C^0(X, M_*) \to C^1(X, M_*)\right). \tag{2.6}$$

Hence, $M_*(X)$ is a subgroup of $C^0(X, M_*)$ and its elements are called *unramified* cycles, see [**26**, p338]. Letting $U \mapsto M_*(U)$ for Zariski opens U, we can turn this into a sheaf $\mathcal{M}_*(U) := M_*(U)$. Moreover, we call

$$C^\bullet(X, M_*) : 0 \longrightarrow M_*(X) \longrightarrow C^0(X, M_*) \xrightarrow{\partial_X} C^1(X, M_*) \xrightarrow{\partial_X} \cdots \tag{2.7}$$

the *Rost complex* of X *with coefficients in* M_*. The first morphism is the subgroup inclusion, all further morphisms are the differential ∂_X. As our scheme is assumed to be n-equidimensional, the sets $X^{(i)}$ for $i > n$ are empty, so $C^i(X, M_*) = 0$, showing that the Rost complexes are two-sided bounded.

2.3. Examples of Cycle Modules. Let us list the most prominent examples of cycle modules and note that they are available for *all* fields F/k (as opposed to the treatise of Rost, who only considers fields F which are finitely generated over k):

(1) The simplest cycle module is given by the Milnor K-ring itself: $M_*(F) := K_*^M(F)$.

(2) Algebraic (Quillen) K-groups also provide an example: $M_*(F) := K_*(F)$. See Rost [**26**, Rmk. 1.12] for details.

(3) The Galois cohomology of certain Galois modules also provides examples. See Rost [**26**, Rmk. 1.11] for details. For this suppose k is a field, $n \geq 1$ an integer coprime to $\operatorname{char} k$ (it is important here that the residue fields of k with respect to *any* discrete valuation in our sense will have the same characteristic because we demand valuations to be trivial on k, so n will still be coprime to it). Suppose $X \to \operatorname{Spec} k$ is smooth. We write $G_F := \operatorname{Gal}(F^{\mathrm{sep}}/F)$ as a short-hand. Let Λ be a *finite n-torsion* Galois module over k, i.e. a continuous G_k-module which is finitely generated as a $\mathbf{Z}/n$-module. Write $\Lambda(q) := \Lambda \otimes \mu_n^{\otimes q}$, $\Lambda(-q) := \operatorname{Hom}(\mu_n^{\otimes q}, \Lambda)$ for $q \geq 0$. In his setup, Rost shows:

Proposition 1 ([**26**, Rmk 1.11, 2.5])**.** *If Λ is a finite n-torsion G_k-module, $(n, \operatorname{char} k) = 1$, the object function*

$$M_*(F) := \coprod_{q \geq 0} H^q(F, \Lambda(q)) \tag{2.8}$$

is a cycle module.

Proof. See [**26**, Rmk 1.11, 2.5]. □

(4) *A new aspect:* For higher local class field theory so-called 'topological Milnor K groups' K_*^{top} are more convenient than ordinary Milnor K groups. Topological Milnor K-theory, see [**9**], [**32**], is useful in order to reflect the continuity features of completed fields, e.g. $K_1^M(\mathbf{Q}_p) = \mathbf{Q}_p^\times$ has a natural topology and there are groups K_*^{top} which reflect such topological features. Moreover, the local reciprocity map is injective on these groups, whereas it has a divisible subgroup as its kernel for ordinary Milnor K-groups. As proven in [**9**], if we disregard the topology, we can at least define algebraically

$$K_n^t(F) := K_n^M(F) / K_n^M(F)_{\mathrm{div}}, \tag{2.9}$$

where $K_n^M(F)_{\mathrm{div}}$ denotes the subgroup of $K_n^M(F)$ of elements of infinite height (= p-divisible elements for all p; for higher local fields this coincides with the maximal divisible subgroup, see [**9**]). This definition makes sense for all fields and has been investigated by Izhboldin, see [**16**, §2.5]. If F has characteristic zero, then unlike ordinary Milnor K groups these groups vanish if n exceeds the cohomological dimension[3] of the field F. We wish to emphasize that the cycle module structure of K_*^M induces all maps **D1**-**D4** to eq. 2.9, in particular K_n^t is a cycle module - and thus our idèle resolution applies to K_n^t-coefficients as well. If L/F are higher local fields, Fesenko has shown that res_F^L / cor_F^L are continuous maps, [**9**, §4.8] (given the definition of the corestriction this is not at all obvious).

We denote the corresponding Zariski sheaf (à la eq. 2.6) by $\mathcal{K}_*^t$. Its cohomology can differ from the one of $\mathcal{K}_*^M$, e.g. naïvely

$$H^0(\operatorname{Spec} \mathbf{C}, \mathcal{K}_1^M) = \mathbf{C}^\times, \qquad \text{whereas} \qquad H^0(\operatorname{Spec} \mathbf{C}, \mathcal{K}_1^t) = 0,$$

while other properties remain generally unaffected, e.g. the analogue of the Bloch-Quillen formula $H^p(X, \mathcal{K}_p) = \mathrm{CH}^p(X)$ holds:

Remark 1. *If X is integral smooth of finite type over a field k then $H^p(X, \mathcal{K}_p^t) \cong \mathrm{CH}^p(X)$.*

Proof. According to the Gersten resolution of the cycle module sheaf on the left-hand side, the group $H^p(X, \mathcal{K}_p^t)$ is the cokernel of

$$\partial_y^x : \coprod\nolimits_{x \in X^{(p-1)}} \kappa(x)^\times / \kappa(x)^\times_{\mathrm{div}} \longrightarrow \coprod\nolimits_{y \in X^{(p)}} \mathbf{Z}/\mathbf{Z}_{\mathrm{div}},$$

---

[3] In characteristic $p > 0$ the p-cohomological dimension is always ≤ 1, whereas the mod p quotient of Milnor K groups can be non-trivial. To obtain a consistent picture, one needs to modify Galois cohomology, see [**17**, §5.1]. We will not go into this.

but $\mathbf{Z}$ has no non-trivial divisible elements (i.e. $\mathbf{Z}_{\mathrm{div}} = 0$) and so the cokernel is the same as for $\mathcal{K}_p^M$. □

We believe it is very interesting to investigate the cycle module K_*^t for the following reason: For higher local fields F one has $K_n^t(F) = K_n^{\mathrm{top}}(F)$ algebraically, but the latter groups are sequential-topological groups. The group operation is sequentially continuous; sadly it is usually *not* continuous, so one does not have a topological group structure (see [**9**]). Still, using such sequential topological structures it might then be possible to equip the whole idèle complex with a well-behaved topological structure. Such a construction could be useful to obtain a theory of 'higher harmonic analysis' on the idèles which might shed new light on L-functions, see §9 for an outline of this circle of ideas.

2.4. Relation to Coniveau Spectral Sequences. In the special case of Galois cohomology coefficients, i.e. as in Prop. 1, the Rost complex appears as the E_1-page of the Bloch-Ogus spectral sequence converging to étale cohomology filtered by coniveau [**5**]. Define a Zariski sheaf $\mathcal{H}^q(\bullet, r)$ as follows

$$\mathcal{H}^q(U,r) := (U \mapsto H^q(U_{\acute{e}t}, \Lambda(q+r))),$$

where U is a Zariski open in X, Λ as in Prop. 1. This is defined in [**5**, beginning of §4]. Note that this is a *Zariski* sheaf. For example, $H^p(\operatorname{Spec} k, \mathcal{H}^q(\bullet, r)) = 0$ for $n \geq 1$ for any field k, quite unlike $H^q((\operatorname{Spec} k)_{\acute{e}t}, \Lambda(q+r))$ which can be non-zero for arbitrarily large q if the field k has sufficiently large Galois cohomological dimension.

Lemma 1. *$\mathcal{H}^q(\bullet, r) = \mathcal{M}_q^{\Lambda(r-q)}$, where $\mathcal{M}_*^{\Lambda(r-q)}$ denotes the Zariski sheaf of the cycle module defined in eq. 2.8 for the Galois module $\Lambda(r)$.*

Proof. This is a rephrasing of [**5**, Thm. 4.2, part (4.2.2)]. □

Proposition 2 (Bloch, Ogus [**5**, Prop. 3.9])**.** *Under these assumptions, there is a spectral sequence*

$$E_1^{p,q} = C^p(X, M_q^{\Lambda(r-q)}) \Longrightarrow N^\bullet H^{p+q}(X_{\acute{e}t}, \Lambda(r)), \tag{2.10}$$

where

(1) *N denotes the coniveau filtration as in*

$$N^p H^q(X_{\acute{e}t}, \mathcal{F}) = \operatorname{im}(\varinjlim_{Z \in Z^p} H^q_Z(X_{\acute{e}t}, \mathcal{F}) \to H^q(X_{\acute{e}t}, \mathcal{F}));$$

(2) *$C^p(X, M_*)$ denotes an entry of the Rost complex as in eq. 2.4;*
(3) *$M_q^{\Lambda(r)}$ is the cycle module as in eq. 2.8 for the Galois module $\Lambda(r)$, $r \in \mathbf{Z}$.*

For the algebraic K-theory cycle module $M_*(F) = K_*(F)$ the Rost complex gives the E_1-page of the Brown-Gersten-Quillen spectral sequence converging to algebraic K-theory filtered by coniveau, see [**25**, Thm. 5.4].

2.5. Pullback & Pushforward. Suppose X, Y are equidimensional excellent schemes (say n_X, n_Y-equidimensional).

2.5.1. *Pushforward.* If X, Y are finite type over a field and $f : X \to Y$ is proper, Rost constructs a pushforward $f_* : C^\bullet(X, M_*) \to C^\bullet(Y, M_*)$, [**26**, §3.4]. This construction carries over to the case of X, Y just equidimensional excellent and f a finite morphism, see Lemma 3 below.

2.5.2. *Pullback.* If X, Y are finite type over a field and $f : X \to Y$ is flat, Rost constructs a pullback operation $f^* : C^\bullet(Y, M_*) \to C^\bullet(X, M_*)$, see [**26**, §3.5] (he also gives a more general pullback operation, for which we have no use). To prolong this to excellent schemes, we say that a *flat morphism of constant relative dimension* r is a flat morphism $f : X \to Y$ such that there is some $r \in \mathbf{Z}$ such that for all $y \in Y$ and all generic points $x \in f^{-1}(\overline{\{y\}})^{(0)}$ one has

(1) $\dim_X x = \dim_Y y + r$.
(2) or equivalently $\dim \mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,y}} \kappa(y) = (n_X - n_Y) - r \geq 0$.

Example 1. *The morphism* $\operatorname{Spec} k[t] \to \operatorname{Spec} k$ *is flat of constant relative dimension* 1.

Example 2 (Example not of finite type)**.** *Let* $(\mathcal{O}, \mathfrak{m})$ *be a regular* 2*-dimensional local domain containing a field and* K *its field of fractions. Then the composition*

$$\operatorname{Spec} \widehat{\mathcal{O}}_{\mathfrak{m}} \otimes_{\mathcal{O}} K \longrightarrow \operatorname{Spec} \widehat{\mathcal{O}}_{\mathfrak{m}} \longrightarrow \operatorname{Spec} \mathcal{O}$$

is flat of constant relative dimension -1*. To see this, note that the scheme on the left is the fiber of the closed point* (0) *in* $\mathcal{O}$*, so* $y = (0)$ *is the only point in the image;* $\widehat{\mathcal{O}}_{\mathfrak{m}} \otimes_{\mathcal{O}} K$ *is a* 1*-dimensional domain (we shall prove this in Lemma 7), so its zero ideal* $x = (0)$ *is the unique generic point of the fiber and so* $r = -1$.

We shall need the following easy fact:

Lemma 2. *Suppose* R *is a universally catenary Noetherian local domain. Then* $f : \operatorname{Spec} \widehat{R} \to \operatorname{Spec} R$ *is flat of constant relative dimension* 0.

For any $y \in Y$ and $x \in X_y^{(0)}$ Rost defines a *multiplicity* $[\mathcal{O}_X, f]_y^x \in \mathbf{Z}$ and a flat pullback

$$f^* := [\mathcal{O}_X, f, r] : C^p(Y, M_*) \longrightarrow C^{p+r}(X, M_*) \qquad \text{(see [\textbf{26}, §3.5])}.$$

We shall use that certain properties of pushforwards and pullbacks carry over to the case of finite pushforwards and flat pullbacks of constant relative dimension.

Lemma 3. *Suppose*

- X, Y *are equidimensional excellent schemes,*
- $f : X \to Y$ *finite,*
- $g : Z \to Y$ *is flat of constant relative dimension* r*,*
- $W := Z \times_Y X$*,*
- f', g' *are the morphisms induced by base change.*

Then the diagrams

$$\begin{tikzcd}
C^p(X, M_*) \arrow[r, "\partial_X"] \arrow[d, "f_*"'] & C^{p+1}(X, M_*) \arrow[d, "f_*"] \\
C^p(Y, M_*) \arrow[r, "\partial_Y"'] & C^{p+1}(Y, M_*)
\end{tikzcd}
\qquad
\begin{tikzcd}
C^p(X, M_*) \arrow[r, "g'^*"] \arrow[d, "f_*"'] & C^{p+r}(W, M_*) \arrow[d, "f'_*"] \\
C^p(Y, M_*) \arrow[r, "g^*"'] & C^{p+r}(Z, M_*)
\end{tikzcd}$$

$$\begin{tikzcd}
C^p(Y, M_*) \arrow[r, "\partial_Y"] \arrow[d, "f^*"'] & C^{p+1}(Y, M_*) \arrow[d, "f^*"] \\
C^{p+r}(Z, M_*) \arrow[r, "\partial_Z"'] & C^{p+r+1}(Z, M_*)
\end{tikzcd}$$

commute. Moreover, $(f_1 \circ f_2)_* = f_{1*} \circ f_{2*}$ *and* $(g_1 \circ g_2)^* = g_2^* \circ g_1^*$ *for* f_1, f_2 *two finite morphisms* $X \overset{f_2}{\to} Y \overset{f_1}{\to} Y'$, g_1, g_2 *two flat constant rel. dim. morphisms* $Z \overset{g_2}{\to} Y \overset{g_1}{\to} Y'$, $r(g_1 \circ g_2) = r(g_1) + r(g_2)$ *for the constant rel. dimensions respectively.*

See [**26**, Prop. 4.1, 4.6], the proofs carry over verbatim (but note that our f is finite as opposed to just being proper). See also [**8**, §49] for the case $M_* = K_*^M$.

Note that some additional care is required when dealing with rings which are not of finite type over a field: Even when assuming excellence, open subschemes might have dimension strictly less than the original scheme; also note that already very basic rings like $k[[t]][s]$ already fail to be equidimensional.

We shall need the following easy facts:

Lemma 4. *Let* $(R, \mathfrak{m})$ *be an* n*-dimensional equicharacteristic complete Noetherian local domain and* $f \in \mathfrak{m}$ *some non-zero element (in particular* $n \geq 1$*). Then* $S := R[f^{-1}]$ *is an* $(n-1)$*-equidimensional domain.*

If I is an ideal of a ring R, the sets $I[[t]] = \{\sum a_i t^i,\ a_i \in I\}$ are ideals in the ring $R[[t]]$. It is easy to see that $R[[t]]/I[[t]] \cong (R/I)\,[[t]]$.

Lemma 5. *Let* $(R, \mathfrak{m})$ *be an* n*-dimensional equicharacteristic complete Noetherian local domain. Then the morphism* $\operatorname{Spec} R[[t]][t^{-1}] \to \operatorname{Spec} R$ *is faithfully flat of constant relative dimension* 0.

Example 3. *For* $n = 1$ *and* $(R, \mathfrak{m}) := (k, (0))$ *a field, one obtains the straightforward example* $\operatorname{Spec} k((t)) \to \operatorname{Spec} k$.

Proof. The morphism is the composition of

$$\operatorname{Spec} R[[t]][t^{-1}] \xrightarrow{g} \operatorname{Spec} R[[t]] \xrightarrow{f} \operatorname{Spec} R.$$

(1) Firstly, f is flat. It is surjective since for every prime y ($\neq R$) in R the prime $y[[t]]$ in $R[[t]]$ is a preimage, so f is faithfully flat. The localization g is flat, but not surjective (the prime (t) does not lie in the image). However, every prime of the shape $y[[t]]$ lies in the image: use that the primes in $R[[t]][t^{-1}]$ correspond in order-preserving bijection to the primes of $R[[t]]$ not containing the element t. For y a prime in R, $R[[t]]/y[[t]] \cong (R/y)\,[[t]]$, so $y[[t]]$ is a prime, and $t \in y[[t]]$ implies $1 \in y$, which is impossible. It follows that the composition $f \circ g$ is surjective, so $f \circ g$ is a faithfully flat morphism.

(2) Fix a prime y in R. Then $\overline{\{y\}} = \operatorname{Spec} R/y$ and since $f \circ g$ is faithfully flat, by Lying-Over the minimal primes over $yR[[t]][t^{-1}]$ are the generic points of the preimage. As $R[[t]]/y \cong (R/y)\,[[t]]$, so that all primes of the shape $y[[t]]$ ($\not\ni t$ as $1 \notin y$) correspond bijectively to the primes $y[[t]] \cdot R[[t]][t^{-1}]$, we see that $x := yR[[t]][t^{-1}]$ is in fact the unique minimal prime in the preimage. Then

$$\begin{aligned} \dim x &= \dim R[[t]][t^{-1}]/x = \dim R[[t]][t^{-1}]/yR[[t]][t^{-1}] \\ &= \dim (R/y)\,[[t]][t^{-1}] \overset{*}{=} \dim R/y = \dim y, \end{aligned}$$

where '$\overset{*}{=}$' follows from Lemma 4. □

2.6. The Gersten Property. We shall need the following general Gersten-conjecture-type property for cycle modules:

Proposition 3 ([**26**, Thm. 6.1])**.** *Suppose* X *is a local ring of a smooth scheme* $X' \to \operatorname{Spec} k$ *and let* M_* *be a cycle module. Then the complex* $C^\bullet(X, M_*)$ *is exact.*

This shows that the sheafification of eq. 2.7 provides a flasque resolution of $\mathcal{M}_*$, [**26**, Cor. 6.5]. Moreover, we need the analogous statement for complete rings. Such a statement is not proven in [**26**], but the general principle of proof is well-known since Quillen's paper on algebraic K-theory and readily applies to cycle modules as well:

Proposition 4. *Suppose* $X \to \operatorname{Spec} k$ *is an equicharacteristic complete regular local scheme with residue field in* $\mathcal{F}$ *and let* M_* *be a cycle module. Then the complex* $C^{\bullet}(X, M_*)$ *is exact.*

We shall use the following lemma, which is very well-known since in similar shape it appears in most proofs of Noether Normalization:

Lemma 6 (Nagata)**.** *Suppose* $f \in F[[w_1, \ldots, w_n]]$ *is non-zero. Then there is a ring automorphism* σ *(fixing* F*) of the shape*

$$w_n \mapsto w_n \qquad w_i \mapsto w_i + w_n^{c_i}$$

for suitable $c_i \in \mathbf{Z}_{\geq 0}$ *and* $i = 1, \ldots, n-1$ *such that* $\sigma f \in F[[w_1, \ldots, w_n]]$ *is* w_n*-distinguished (in the sense of Weierstraß).*

Proof of Prop. 4. (Adaption of [**25**, Thm. 5.13]) Write $X =: \operatorname{Spec} R$ with $(R, \mathfrak{m})$ a complete regular local domain as in the claim. The injectivity $M_*(X) \hookrightarrow C^0(X, M_*)$ follows from the very definition of $M_*(X)$ as a kernel, so we only need to show that given some $\alpha = (\alpha_y)_{y \in X^{(p+1)}}$ as in the middle term of

$$\cdots \longrightarrow \coprod_{x \in X^{(p)}} M_*(\kappa(x)) \xrightarrow{\partial_X} \coprod_{y \in X^{(p+1)}} M_{*-1}(\kappa(y)) \xrightarrow{\partial_X} \coprod_{z \in X^{(p+2)}} M_{*-2}(\kappa(z))$$

such that $\partial_X \alpha = 0$, there exists some $\beta = (\beta_x)_{x \in X^{(p)}}$ with $\partial_X \beta = \alpha$. Since R is assumed regular and therefore all its localizations are regular and thus factorial, for every local ring $\mathcal{O}_{X,x}$ the codimension 1 closed subschemes $\overline{\{y\}}_{\mathcal{O}_{X,x}}$ (this means we take the Zariski closure in $\operatorname{Spec} \mathcal{O}_{X,x}$) are principal divisors and cut out by some irreducible elements, say $\pi_y \in \mathcal{O}_{X,x}$. Then $\overline{\{y\}}_{\mathcal{O}_{X,x}} = \operatorname{Spec} \mathcal{O}_{X,x}/(\pi_y)$. Assume $X = \operatorname{Spec} \mathcal{O}_{X,x}$ for the moment. Next, by Cohen's Structure Theorem there is a (non-canonical) ring isomorphism

$$\mathcal{O}_{X,x} \simeq \kappa(x)[[w_1, \ldots, w_p]]$$

since $\operatorname{ht}(x) = p$. After possibly changing the isomorphism by Lemma 6 we can assume that π_y is distinguished, i.e. the w_p-coefficients of π_y do not all lie in the ideal $(w_1, \ldots, w_{p-1})$. Then by Weierstraß Preparation there is a unit $u \in \mathcal{O}_{X,x}^{\times}$ and a Weierstraß polynomial $f \in \kappa(x)[[w_1, \ldots, w_{p-1}]][\tilde{w}_p]$ such that $\pi_y = u \cdot f$. Hence, w.l.o.g. $\pi_y = f$ as this generates the same ideal. Next, we mimick the diagram of [**26**, *proof of* Prop. 6.4.]

$$\begin{tikzcd}
 & Z \arrow[dl, "g"] \arrow[dr, "\pi"] & \\
Y \arrow[ur, "\sigma"] \arrow[rr, "i"] \arrow[dr] & & X \arrow[dl] \\
 & A, &
\end{tikzcd}$$

where in our setup

- $Y := \operatorname{Spec} \mathcal{O}_{X,x} / (\pi_y)$;
- $X := \operatorname{Spec} \mathcal{O}_{X,x} = \operatorname{Spec} \kappa(x)[[w_1, \ldots, w_p]]$;
- $A := \operatorname{Spec} \kappa(x)[[w_1, \ldots, w_{p-1}]]$;
- $Z := Y \times_A X$.

Here the arrows g, π are the product projections. The arrow $X \to A$ is not smooth (unlike its counterpart in [**26**, *proof of* Prop. 6.4.]), but flat of constant relative dimension 1. The arrow $Y \to A$ is a finite morphism since by our Weierstraß Preparation above we have

$$\mathcal{O}_Y = \mathcal{O}_{X,x} / (\pi_y) = \kappa(x)[[w_1, \ldots, w_{p-1}]][\tilde{w}_p] / (f(\tilde{w}_p)).$$

The arrow $i : Y \to X$ is a closed immersion, so it is finite. Finally, note that

$$\begin{aligned} \mathcal{O}_Z &= \kappa(x)[[w_1, \ldots, w_{p-1}]][\tilde{w}_p] / (f(\tilde{w}_p)) \ldots \\ &\qquad \ldots \otimes_{\kappa(x)[[w_1, \ldots, w_{p-1}]]} \kappa(x)[[w_1, \ldots, w_{p-1}]][[w_p]] \\ &= \kappa(x)[[w_1, \ldots, w_{p-1}, w_p]][\tilde{w}_p] / (f(\tilde{w}_p)), \end{aligned} \tag{2.11}$$

a computation which crucially uses that $Y \to A$ is finite. This immediately gives us the closed immersion σ coming from the ideal (w_p) in $\mathcal{O}_Z$. This realizes Y as the underlying integral closed subscheme of Z of a rationally trivial Weil divisor. The remaining argument proceeds as in the proof of [**26**, Prop. 6.4]: We let V be the open complement of the closed subset $\sigma(Y)$ in Z so that $Z = V \cup \sigma(Y)$ disjointly. Algebraically,

$$\mathcal{O}_V = \kappa(x)[[w_1, \ldots, w_{p-1}, w_p]][w_p^{-1}][\tilde{w}_p] / (f(\tilde{w}_p)).$$

(*Beware:* This is *not* isomorphic to $\kappa(x)((w_p))[[w_1, \ldots, w_{p-1}]][\tilde{w}_p] / (f(\tilde{w}_p))$ - in the latter ring elements may have w_p-degrees which are unbounded from below. This is not possible in the ring underlying V.) We have $\dim V = p - 1$ while $\dim Z = p$, i.e. the open subscheme V has strictly lower dimension than Z (to see this use that $\mathcal{O}_V$ is an integral extension of a ring as in Lemma 4). Let $g' : V \to Y$ be the restriction of the projection $Z \to Y$ to the open subscheme V. This morphism is flat of constant relative dimension 0 by Lemma 5 (whereas the full $g : Z \to Y$ has constant relative dimension 1). Define

$$H := \pi_* \circ j_* \circ \{w_p\} \circ g'^* \qquad \text{(not to be confused with } \tilde{w}_p\text{)}$$

$$C^p(Y, M_*) \to C^p(V, M_*) \to C^p(V, M_{*+1}) \to C^p(Z, M_{*+1}) \to C^p(X, M_{*+1}),$$

where

- g'^* is the flat pullback of constant rel. dim. 0 (Lemma 5);
- $\{w_p\}$ denotes the left-multiplication by units on $C^\bullet$, see [**26**, §3.6] for the definition of this operation, but essentially this just uses the left-K_1^M-module operation **D3** on cycle modules, and note that $w_p \in \mathcal{O}_V^\times$ is indeed a unit on V (in fact we just chose V as to make this statement valid);
- $j : V \to Z$ is the open immersion (and flat of constant relative dimension 0) and j_* is a *non*-proper pushforward along this open immersion. *Beware:* As j is not proper, $\partial \circ j_* \neq j_* \circ \partial$.
- $\pi : Z \to X$ is the product projection. It is a *finite* morphism, one sees this by direct inspection of eq. 2.11 or abstractly by base change from the finiteness of $Y \to A$.

As for the proof of [**26**, Prop. 6.4] one computes that H is a chain homotopy between the closed immersion pushforward i_* and the zero morphism:

$$\partial_X \circ H + H \circ \partial_Y = i_* - 0 : C^p(Y, M_*) \to C^{p+1}(X, M_{*+1}).$$

This completes the proof. $\square$

3. Preparations

3.1. 2-dimensional Local Rings. Suppose $(\mathcal{O}, x)$ is a smooth 2-dimensional local domain with residue field $\kappa(x)$ and essentially of finite type over a field k so that $\kappa(x)/k$ is a finite field extension. We write $X := \operatorname{Spec} \mathcal{O}$ for simplicity.

3.2. Notation. Let us agree on the following notation:

- $\widehat{\mathcal{O}}_x$ is the x-adic completion of $\mathcal{O}$;
  This is a 2-dimensional complete regular local domain;
- $\widehat{\mathcal{O}}_y$, for every height 1 prime $y \in (\operatorname{Spec} \mathcal{O})^{(1)}$, is the y-adic completion of the localization $\mathcal{O}_y$;
  This is a 1-dimensional complete regular local domain. Hence, it is a CDVR with residue field $\kappa(y)$;
- $\widehat{\mathcal{O}}_{x,y'_i}$, for every height 1 prime $y \in (\operatorname{Spec} \mathcal{O})^{(1)}$ and y'_i a prime in $\widehat{\mathcal{O}}_x$ over y (see Lemma 7.(4) for details on this), is the y'_i-adic completion of the localization of $\widehat{\mathcal{O}}_x$ at y'_i, one could also write $\widehat{(\widehat{\mathcal{O}}_x)_{y'_i}}$, which is a clearer, but really unwieldy notation;
  This ring is a 1-dimensional complete regular local domain. Thus, it is a CDVR and the residue field is $\kappa(y'_i)$ of $\operatorname{Spec} \widehat{\mathcal{O}}_x$;
- $K := \operatorname{Frac} \mathcal{O}$ is the field of fractions of $\mathcal{O}$;
- $\widehat{K}_y := \operatorname{Frac} \widehat{\mathcal{O}}_y$ is the field of fractions of $\widehat{\mathcal{O}}_y$;
  This is a 1-dimensional local field[4] with residue field $\kappa(y)$;
- $\widehat{K}_x := \operatorname{Frac} \widehat{\mathcal{O}}_x$ (this does *not* appear in the idèles, rather $\widehat{\mathcal{O}}_x \otimes K$ will need to be used)
- $\widehat{K}_{x,y'_i} := \operatorname{Frac} \widehat{\mathcal{O}}_{x,y'_i}$ is the field of fractions of $\widehat{\mathcal{O}}_{x,y'_i}$;
  This is a 2-dimensional local field with residue fields $\kappa(y'_i)$ and $\kappa(x)$:

$$\begin{tikzcd}
\widehat{K}_{x,y'_i} & & \\
\widehat{\mathcal{O}}_{x,y'_i} \arrow[u] \arrow[r] & \kappa(y'_i) & \\
& (\widehat{\mathcal{O}}_x/y'_i) \arrow[u] \arrow[r] & \kappa(x).
\end{tikzcd} \tag{3.1}$$

  Here the upward arrows denote the inclusion of the rings of integers (= the valuation rings) into their fields of fractions and the rightward arrows denote the surjections on the relative residue fields. See [**19**] for an overview regarding higher local fields.

---

[4] for us this means: a complete discrete valuation field. We do not demand the residue field to be perfect or (quasi-)finite as usual.

3.3. Points on these local schemes. We shall need a few facts about the structure of these rings: The natural morphism $j : \operatorname{Spec}\widehat{\mathcal{O}}_x \to \operatorname{Spec}\mathcal{O}$ is faithfully flat and thus surjective and we decompose

$$\operatorname{Spec}\widehat{\mathcal{O}}_x = \widehat{X}^{\langle 0\rangle} \cup \widehat{X}^{\langle 1\rangle} \cup \widehat{X}^{\langle 2\rangle}$$

disjointly, where $\widehat{X}^{\langle i\rangle}$ denotes the set of primes $\widehat{y}$ such that $j(\widehat{y}) \in (\operatorname{Spec}\mathcal{O})^{(i)}$, i.e. it contains the primes whose contractions to $\mathcal{O}$ are of height $0 \leq i \leq 2$. The primes itself may have a *different* height in $\widehat{\mathcal{O}}_x$, we clarify this now:

Definition 2. *We shall call the height* 1 *primes* $\widehat{y}$ *of* $\widehat{\mathcal{O}}_x$ *such that* $\widehat{y} \notin X^{\langle 1\rangle}$ transcendental *height* 1 *primes.*

Lemma 7. *We have:*

1. $\widehat{X}^{\langle 2\rangle} = \{x\widehat{\mathcal{O}}_x\}$. *This is the single height* 2 *prime in* $\widehat{\mathcal{O}}_x$.
2. *The ring* $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ *is a* 1*-dimensional Noetherian domain. In fact, it is a principal ideal domain.*
3. $i : \widehat{X}^{\langle 0\rangle} \overset{\cong}{\to} \operatorname{Spec}\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ *(as sets, i.e. there is a bijection). This bijection preserves residue fields, i.e.*

$$\kappa_{\widehat{\mathcal{O}}_x}(\widehat{y}) \cong \kappa_{\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K}(i\widehat{y}) \qquad \textit{(where } \kappa_R(x) = \mathcal{O}_{R,x}/\mathfrak{m}_{R,x}\textit{)}$$

for all $\widehat{y} \in \widehat{X}^{\langle 0\rangle}$. *Moreover,*

$$\widehat{X}^{\langle 0\rangle} = X^{(1)}_{\text{transcendental}} \cup \{(0)\}.$$

4. *For every* $y \in X^{(1)}$ *the preimage* $j^{-1}(y) = \{\widehat{y}_1, \ldots, \widehat{y}_r\}$ *is finite with* $\widehat{y}_i \in \widehat{X}^{\langle 1\rangle}$ *and every element in* $\widehat{X}^{\langle 1\rangle}$ *arises this way.*

Proof. **(1)** is obvious. **(2)** $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ is a localization of $\widehat{\mathcal{O}}_x$, so it is Noetherian. We may regard it as a subring of the field of fractions $\widehat{K}_x = \operatorname{Frac}\widehat{\mathcal{O}}_x$. Choosing a non-canonical isomorphism $\widehat{\mathcal{O}}_x \simeq \kappa(x)[[s,t]]$ by Cohen's Structure Theorem, the transcendental height 1 prime $t - s\exp(s)$ proves that $\dim(\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K) \geq 1$. On the other hand, $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ is a localization of $\widehat{\mathcal{O}}_x[t^{-1}]$, and the latter satisfies $\dim\widehat{\mathcal{O}}_x[t^{-1}] = 1$ by Lemma 4. This shows $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ is 1-dimensional. Moreover, as $\widehat{\mathcal{O}}_x$ is factorial, $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ must also be factorial, so the height 1 primes are principal, so the ring must be a PID.

(3) For $\widehat{y} \in \widehat{X}^{\langle 0\rangle}$, $j(\widehat{y})$ is a height zero prime and thus the zero ideal since $\mathcal{O}$ is a domain. Hence, $\widehat{X}^{\langle 0\rangle}$ bijects to the primes in the formal fiber $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$. As we know the structure of the latter scheme by part **(2)** of this lemma, we get the zero ideal plus a lot(!) of height 1 primes. These are precisely the transcendental primes. For the residue field at some $\widehat{y} \in \widehat{X}^{\langle 1\rangle}$ we use that $\widehat{y} \cap K = (0)$ (as $\widehat{y}$ maps to the zero ideal) and thus in terms of localizing a multiplicative subset,

$$\begin{aligned}\kappa_{\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K}(\widehat{y}) &= \operatorname{Frac}((\widehat{\mathcal{O}}_x)_{\mathcal{O}-\{0\}}/\widehat{y}) \\ &= \operatorname{Frac}((\widehat{\mathcal{O}}_x/\widehat{y})_{\overline{\mathcal{O}-\{0\}}}) = \operatorname{Frac}\widehat{\mathcal{O}}_x/\widehat{y} = \kappa_{\widehat{\mathcal{O}}_x}(\widehat{y}),\end{aligned}$$

where $\overline{\mathcal{O}-\{0\}}$ denotes the image in $\widehat{\mathcal{O}}_x/\widehat{y}$.

(4) Let $y \in X^{(1)}$ be given. By definition all primes in the preimage $j^{-1}(y)$ lie in $\widehat{X}^{\langle 1\rangle}$ and conversely the elements in this set map to height 1 primes. We recall that

j is surjective, so it suffices to prove that each preimage $j^{-1}(y)$ is finite. To see this, note that the formal fiber is

$$\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} \kappa(y) = \widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} \operatorname{Frac} \mathcal{O}/y,$$

i.e. a localization of $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} \mathcal{O}/y = \widehat{\mathcal{O}}_x / y\widehat{\mathcal{O}}_x$. However, $\widehat{\mathcal{O}}_x / y\widehat{\mathcal{O}}_x = \widehat{\mathcal{O}/y\mathcal{O}_x}$ and we know that since $\mathcal{O}/y$ is a 1-dimensional domain, the x-adic completion is a reduced x-adically complete local ring with finitely many minimal primes. $\square$

Lemma 8. *There is an exact sequence of graded abelian groups*

$$0 \to M_*(\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K) \to M_*(\widehat{K}_x) \to \coprod_{\widehat{y}} M_{*-1}(\kappa(\widehat{y})) \to 0, \tag{3.2}$$

where $\widehat{y}$ runs through the transcendental height 1 primes of $\widehat{\mathcal{O}}_x$.

If some Gersten-type statement (i.e. a statement like Prop. 3 or Prop. 4) a priori held true for $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$, this would of course readily imply the result since $\widehat{K}_x$ is the field of fractions and the height 1 primes are as said. However, $\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ is *not* a semilocal ring, so a little work needs to be invested.

PROOF. Using Lemma 7 the Rost complex of $\operatorname{Spec} \widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$ is the first row of the diagram below, where $\widehat{y}$ runs through the height 1 (= maximal) primes $\operatorname{Spec} \widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K$, i.e. primes bijecting to the transcendental primes of Def. 2; y' runs through all height 1 primes of $\widehat{\mathcal{O}}_x$:

$$\begin{tikzcd} \cdots \arrow[r] & M_*(\widehat{K}_x) \arrow[r] \arrow[d, equal] & \coprod_{\widehat{y}} M_{*-1}(\kappa(\widehat{y})) \arrow[r] \arrow[d, shift left] & 0 & \\ \cdots \arrow[r] & M_*(\widehat{K}_x) \arrow[r] & \coprod_{y'} M_{*-1}(\kappa(\widehat{y})) \arrow[r] \arrow[u, shift left] & M_{*-2}(\kappa(x)) \arrow[r] & 0. \end{tikzcd} \tag{3.3}$$

Prop. 4 for $\operatorname{Spec} \widehat{\mathcal{O}}_x$ yields exactness of the lower row. Exactness of the upper row on the left (not displayed above) is clear since $M_*(\widehat{\mathcal{O}}_x \otimes_{\mathcal{O}} K)$ is just defined to be the kernel of the map in the middle. The arrows '↑↓' denote the projection/inclusion of direct summands into the other coproduct. Exactness at $M_*(\widehat{K}_x)$ is clear in both rows. To see surjectivity on the right in the upper row, note that in the exact sequence (exact by Prop. 3 for $\operatorname{Spec} \mathcal{O}_x$)

$$\cdots \to M_*(K) \to \coprod\nolimits_{\operatorname{ht} y=1} M_{*-1}(\kappa(y)) \to M_{*-2}(\kappa(x)) \to 0 \tag{3.4}$$

surjects onto $M_{*-2}(\kappa(x))$ just as the lower row of eq. 3.3 does. However, in eq. 3.4 y runs only through non-transcendental primes, so under the arrow '↑' these summands are sent to zero anyway. Thus, for any element on the right in the upper row of eq. 3.3, we can find a candidate in the lower row being sent to zero in $M_{*-2}(\kappa(x))$, then use exactness of the lower row. $\square$

4. Two-dimensional Idèle Sheaves with Cycle Module Coefficients

Overview: In this section we define the sheaves which give the idèles for surfaces. We fix an integral smooth surface X over a field k, η denotes its generic point and pick a cycle module M_* over k.
Notation: We shall use a simplified notation to define these sheaves. We shall write expressions like

$$\prod\nolimits_y M_*(\widehat{K}_y) \quad \text{as a shorthand for} \quad F : U \to \prod_{y \in U^{(1)}} M_*(\widehat{K}_y), \tag{4.1}$$

and the latter is easily seen to be a flasque sheaf (the restriction to smaller opens is the identity on factors). Analogously for $\prod_x$, where x runs through all closed points $U^{(2)}$, i.e. $U \to \prod_{x \in U^{(2)}} M_*(\widehat{K}_x)$. Writing $\prod_x \coprod_y$ or $\prod_{x,y'_i}$, the second parameter y (or y'_i, y', $\widehat{y}$) runs through the height 1 primes of $\mathcal{O}_x$ (or non-transcendental height 1 in $\widehat{\mathcal{O}}_x$, any height 1 in $\widehat{\mathcal{O}}_x$, or transcendental height 1 in $\widehat{\mathcal{O}}_x$ respectively). For y this is equivalent to y running though $U^{(1)}$ with $x \in \overline{\{y\}}$ ('integral curves passing through the closed point x'). For y'_i it is equivalent to running through the primes of the fibers of $\operatorname{Spec} \widehat{\mathcal{O}}_x \to \operatorname{Spec} \mathcal{O}$ over primes $y \in U^{(1)}$. For y' it is equivalent to also incorporating the closed points in the fiber of (0), $\widehat{y}$ is just the transcendental primes.

We shall also need certain finiteness conditions constraining the sheaves of eq. 4.1:

(1) The prime superscript ’′’ in $\prod'_y M_*(\widehat{K}_y)$ means that we consider only those elements $(\alpha_y)_y$ such that

$$\alpha_y \in M_*(\widehat{\mathcal{O}}_y) \text{ for all but finitely many } y.$$

(2) The double prime superscript ’′′’ in $\prod''_x M_*(\widehat{K}_x)$ means that we consider only those elements $(\alpha_x)_x$ such that

$$\alpha_x \in M_*(\widehat{\mathcal{O}}_x \otimes K_\eta) \text{ for all but finitely many } x.$$

(3a) The triple prime superscript ’′′′’ in $\prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i})$ means that we consider only those elements $(\alpha_{x,y'_i})_{x,y'_i}$ such that

- **(a):** $\forall x : \alpha_{x,y'_i} \in M_*(\widehat{\mathcal{O}}_{x,y'_i})$ for all but finitely many y'_i;
- **(b):** $\sum_{y'_i} \partial_x^{y'_i} \partial_{y'_i}^{\widehat{K}_{x,y'_i}} (\alpha_{x,y'_i}) = 0 \in M_{*-2}(\kappa(x))$ for all but finitely many x.

The condition **(a)** ensures that the sum in **(b)** is finite.

Example 4. *Note that (a) still allows $\alpha_{x,y'_i} \notin M_*(\widehat{\mathcal{O}}_{x,y'_i})$ for infinitely many primes y'_i as shown in this example:*



The lines represent primes y'_i with $\partial_{y'_i} \alpha_{x,y'_i} \neq 0$. In this particular example infinitely many closed points x are involved, so the condition (b) is non-trivial.

(3b) The superscript ’′′′’ in $\prod'''_x \coprod_{y'_i} M_{*-1}(\kappa(y'_i))$ means that

- **(a):** y' just runs through non-transcendental primes $y' = y'_i$ for all but fin. many x. ($\Leftrightarrow$ there are only finitely many x where the coproduct $\coprod_{y'} M_{*-1}(\kappa(y'))$ has non-trivial contribution at some transcendental prime $y' = \widehat{y}$)
- **(b):** $\sum_{y'_i} \partial_x^{y'_i} \alpha_{x,y'_i} = 0 \in M_{*-2}(\kappa(x))$ for all but finitely many x.

The sum over non-transcendental primes y'_i in **(b)** is automatically finite.

Construction of Idèle Sheaves: We define a tricomplex W_0 of sheaves on X with values in abelian groups by

$$\begin{tikzcd}
 & \prod''_x M_*(\widehat{K}_x) \arrow[dl] & & \prod''_x M_*(\widehat{K}_x) \arrow[ll] \arrow[dl] \\
\prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i}) & & \prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i}) \arrow[ll] & \\
 & \coprod_\eta M_*(K_\eta) \arrow[uu] \arrow[dl] & & \coprod_\eta M_*(K_\eta) \arrow[uu] \arrow[ll] \arrow[dl] \\
\prod'_y M_*(\widehat{K}_y) \arrow[uu] & & \prod'_y M_*(\widehat{K}_y) \arrow[ll] \arrow[uu] &
\end{tikzcd}$$

The arrows are defined by the respective flat pullbacks, e.g. $M_*(K_\eta) \to M_*(\widehat{K}_y)$ is induced from the flat pullback/the restriction $\operatorname{res}^{\widehat{K}_y}_{K_\eta}$ of the cycle module. These sheaves are all flasque (easy to see) and considering the arrows in the direction '$\longleftarrow$' we conclude that the diagonal complex of W_0 is exact (because all these arrows are the identity map).

Lemma 9. *The morphisms in the above diagram are well-defined.*

Proof. For some arrows it may be necessary to convince ourselves that they are well-defined:

(1) $\coprod_\eta M_*(K_\eta) \to \prod'_y M_*(\widehat{K}_y)$ exists since for $\alpha_\eta \in M_*(K_\eta)$ there are only finitely many y with $\partial^\eta_y \alpha_\eta \neq 0$ (this is the cycle module axiom **FD** in [**26**, Def. 2.1], but the latter is equivalent to $\operatorname{res}^{\widehat{K}_y}_{K_\eta} \alpha_\eta \notin M_*(\widehat{\mathcal{O}}_y)$.

(2) $\prod'_y M_*(\widehat{K}_y) \to \prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i})$ exists: Condition (3a).(a) holds by the finiteness condition of $\prod'_y$. For each of the finitely many $\alpha_y \in M_*(\widehat{K}_y)$ with $\partial^{\widehat{K}_y}_y \alpha_y \neq 0$ there are only finitely many x such that $\partial^y_x(\partial^{\widehat{K}_y}_y \alpha_y) \neq 0$, again by **FD**. For each y'_i we have (by axiom **R3a**)

$$\partial^{\widehat{K}_{x,y'_i}}_{y'_i} \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} \alpha_y = \operatorname{res}^{\kappa(y'_i)}_{\kappa(y)} \partial^{\widehat{K}_y}_y \alpha_y,$$

and by applying $\partial^{y'_i}_x$ we deduce that in total there are only finitely many chains (y'_i, x) with $\partial^{y'_i}_x \partial^{\widehat{K}_{x,y'_i}}_{y'_i} \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} \alpha_y \neq 0$ at all. Thus, the sum in (3a).(b) can possibly be non-zero only for finitely many x, which is precisely what we needed to show.

(3) $\prod''_x M_*(\widehat{K}_x) \to \prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i})$ exists: For all but finitely many x we have $\alpha_x \in M_*(\widehat{\mathcal{O}}_x \otimes K_\eta)$; assume we treat a factor x of the latter type: Then $\partial^{\widehat{K}_x}_{\widehat{y}} \alpha_x = 0$ for all transcendental primes $\widehat{y}$ by Lemma 8. Thus, regarding condition (3a).(b) we compute (where y' runs through *all* height 1 primes of $\widehat{\mathcal{O}}_x$)

$$\sum_{y'} \partial^{y'}_x \partial^{\widehat{K}_{x,y'_i}}_{y'} (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_x} \alpha_x) = \sum_{y'} \partial^{y'}_x \operatorname{res}^{\kappa(y')}_{\kappa(y')} \partial^{\widehat{K}_x}_{y'} (\alpha_x) = 0 \tag{4.2}$$

since we have $\partial^2 = 0$ in $M_*(\widehat{K}_x) \to \coprod_{y'} M_{*-1}(\kappa(y')) \to M_{*-2}(\kappa(x))$. The sum in (3a).(b) is just the above sum, but with y' only running over *non-transcendental* primes y'_i. However, since $\alpha_x \in M_*(\widehat{\mathcal{O}}_x \otimes K_\eta)$, we have $\partial^{\widehat{K}_x}_{\widehat{y}}(\alpha_x) = 0$ for all non-transcendental primes, so it is actually the same sum. Hence, it remains to treat

the finitely many remaining x having $\partial_{\widehat{y}}^{\widehat{K}_x}(\alpha_x) \neq 0$ for some transcendental prime $\widehat{y}$. But (3a).(b) only needs to hold for all but finitely many x, so this case does not matter.
For the remaining arrows it is obvious that they are well-defined. □

We define a tricomplex W_1 as follows:

$$\begin{tikzcd}
& \coprod_x \coprod_{\widehat{y}} M_{*-1}(\kappa(\widehat{y})) \arrow[dl] & & \prod'''_x \coprod_{y'} M_{*-1}(\kappa(y')) \arrow[ll] \arrow[dl] \\
0 & & \prod'''_x \coprod_{y'_i} M_{*-1}(\kappa(y'_i)) \arrow[ll] & \\
& 0 \arrow[uu] \arrow[dl] & & \coprod_y M_{*-1}(\kappa(y)) \arrow[ll] \arrow[uu] \arrow[dl] \\
0 \arrow[uu] & & \coprod_y M_{*-1}(\kappa(y)) \arrow[ll] \arrow[uu] &
\end{tikzcd}$$

Again, all these sheaves are flasque. Using the fact that the set of all height 1 primes y' in $\widehat{\mathcal{O}}_x$ decomposes disjointly into the transcendental primes $\widehat{y}$ and non-transcendental primes y'_i, it is easy to see that the diagonal complex of W_1 is also exact.

Lemma 10. *The morphisms in the above diagram are well-defined.*

Proof. Easy. □

Finally, we define a (rather trivial) tricomplex W_2 as follows:

$$\begin{tikzcd}
& 0 \arrow[dl] & & \coprod_x M_{*-2}(\kappa(x)) \arrow[ll] \arrow[dl] \\
0 & & 0 \arrow[ll] & \\
& 0 \arrow[uu] \arrow[dl] & & \coprod_x M_{*-2}(\kappa(x)) \arrow[ll] \arrow[uu] \arrow[dl] \\
0 \arrow[uu] & & 0 \arrow[ll] \arrow[uu] &
\end{tikzcd}$$

Again, the sheaves are flasque and the diagonal complex is exact.

Having given these definitions, we turn this data into a complex with values in tricomplexes of sheaves

$$W_\bullet : 0 \longrightarrow W_0 \xrightarrow{d} W_1 \xrightarrow{d} W_2 \longrightarrow 0,$$

where the differentials d are induced on the factors mostly by the boundary maps ∂_y^η and ∂_x^y in the obvious way (we give the details below). If we agree on the indexing pattern

(4.3)
$$\begin{tikzcd}
& 02 \arrow[dl] & & 2 \arrow[ll] \arrow[dl] \\
012 & & 12 \arrow[ll] & \\
& 0 \arrow[uu, dashed] \arrow[dl] & & X \arrow[ll] \arrow[uu] \arrow[dl] \\
01 \arrow[uu] & & 1 \arrow[ll] \arrow[uu] &
\end{tikzcd}$$

(written as superscripts) for the individual entries of tricomplexes, the cross sections of the complex $W_\bullet$ in the individual entries are as follows:

$$\begin{array}{llcccc}
W_\bullet^X & : 0 \to & \coprod_\eta M_*(K_\eta) & \overset{\partial_y^\eta}{\to} & \coprod_y M_{*-1}(\kappa(y)) & \overset{\partial_x^y}{\to} & \coprod_x M_{*-2}(\kappa(x)) \\
W_\bullet^0 & : 0 \to & \coprod_\eta M_*(K_\eta) & \to & 0 & \to & 0 \\
W_\bullet^1 & : 0 \to & \prod'_y M_*(\widehat{K}_y) & \overset{\partial_y^\eta}{\to} & \coprod_y M_{*-1}(\kappa(y)) & \to & 0 \\
W_\bullet^2 & : 0 \to & \prod''_x M_*(\widehat{K}_x) & \overset{\partial_{y'}^\eta}{\to} & \prod'''_x \coprod_{y'} M_{*-1}(\kappa(y')) & \overset{\partial_x^{y'}}{\to} & \coprod_x M_{*-2}(\kappa(x)) \\
W_\bullet^{01} & : 0 \to & \prod'_y M_*(\widehat{K}_y) & \to & 0 & \to & 0 \\
W_\bullet^{02} & : 0 \to & \prod''_x M_*(\widehat{K}_x) & \overset{\partial_{\widehat{y}}^\eta}{\to} & \coprod_x \coprod_{\widehat{y}} M_{*-1}(\kappa(\widehat{y})) & \to & 0 \\
W_\bullet^{12} & : 0 \to & \prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i}) & \overset{\partial_{y'_i}^\eta}{\to} & \prod'''_x \coprod_{y'_i} M_{*-1}(\kappa(y'_i)) & \to & 0 \\
W_\bullet^{012} & : 0 \to & \prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i}) & \to & 0 & \to & 0
\end{array}$$

where writing ∂_y^x stands for $\sum \partial_y^x$ (see eq. 2.2) over all codimension 1 points $y \in \overline{\{x\}}^{(1)}$. It will soon play a key role that $W_\bullet^X$ is just Rost's Gersten resolution for the Zariski sheaf $\mathcal{M}_*$ of the cycle module M_*, as in eq. 2.7, turned into sheaves. The above cross section complexes are generally not exact (e.g. $W_\bullet^{012}$ or $W_\bullet^0$).

Lemma 11. *The morphisms in the above diagram are well-defined.*

Proof. Just a variation of the proof of Lemma 9. ☐

Moreover, we define sheaves 'idèles with a reciprocity constraint' by

(1) $\prod_y^{\prime+\text{recip.}} M_*(\widehat{K}_y) := \ker\left(\prod'_y M_*(\widehat{K}_y) \overset{\partial_x^y \partial_y}{\to} \coprod_x M_{*-2}(\kappa(x))\right)$, and

(2) $\prod_{x,y'_i}^{\prime\prime\prime+\text{recip.}} M_*(\widehat{K}_{x,y'_i}) := \ker\left(\prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i}) \overset{\partial_x^{y'_i} \partial_{y'_i}}{\to} \coprod_x M_{*-2}(\kappa(x))\right)$.

Remark 2. *Here in (1) the word 'reciprocity constraint' refers to the fact that an element in $M_*(K)$ when diagonally mapped to $\prod'_y M_*(\widehat{K}_y)$ automatically satisfies $\sum_y \partial_x^y \partial_y x = 0$. As this sum runs over all curves y through a fixed closed point x, this is sometimes called a reciprocity law 'around a point'. Thus, for us, by a reciprocity constraint we understand a necessary criterion to be in the image of the diagonal map. Analogously for (2).*

The morphisms we are taking kernels of are well-defined by the finiteness conditions (1) and (3a).(b) respectively. It is easy to see that the second sheaf is flasque (since the product is indexed over $x \in U^{(2)}$). Unlike all other sheaves discussed in this section so far (apart from $\mathcal{M}_*$), the sheaf $\prod_y^{\prime+\text{recip.}} M_*(\widehat{K}_y)$ is usually *not* flasque:

Example 5. *Consider the restriction of affine 2-space $\mathbf{A}_k^2 := \operatorname{Spec} k[s,t]$ with the closed point $x = (s,t)$ to the smaller open $\mathbf{A}_k^2 - x$ and the Milnor K cycle module $M_* = K_*^M$. The element*

$$[(s), \{s,t\}] + \prod_{y \neq (s)} [y, 0] \in \prod_y{}^{\prime} K_2^M(\widehat{K}_y)$$

by $\Sigma \partial_y$ is mapped to $[(s), \{t\}] \in \coprod_y K_1^M(\kappa(y))$, and thus to $1 \in K_0^M(\kappa(x))$. Hence, the element is a section of $\prod_y^{\prime+recip.} M_*(\widehat{K}_y)$ on $\mathbf{A}_k^2 - x$ which does not *lie in the* *image of the restriction of sections from the open* $\mathbf{A}^2_k$. *Note however that the sheaf is flasque on the degree* ≤ 1 *parts since* $K^M_{-1}(F) = K^M_{-2}(F) = 0$ *for all fields* F.

Using the long exact sequence in cohomology and the flasqueness of the two sheaves $\prod'_y M_*(\widehat{K}_y)$ and $\coprod_x M_{*-2}(\kappa(x))$, we can at least conclude

$$H^p(X, \prod_y^{\prime+\text{recip.}} M_*(\widehat{K}_y)) = 0 \qquad \text{for} \qquad p \geq 2.$$

Next, we compute the cohomology of the complex $W_\bullet$ (which is a tricomplex since $W_\bullet$ is a complex of tricomplexes). We get:

Lemma 12. $H^0(W_\bullet)$ *is the tricomplex of sheaves given by*

(4.4)
$$\begin{tikzcd}
 & \prod_x M_*(\widehat{\mathcal{O}}_x \otimes K_\eta) \arrow[dl] & & \prod_x M_*(\widehat{\mathcal{O}}_x), \arrow[ll] \arrow[dl] \\
\prod'''_{x,y'_i} M_*(\widehat{K}_{x,y'_i}) & & \prod_{x,y'_i} M_*(\widehat{\mathcal{O}}_{x,y'_i}) \arrow[ll] & \\
 & \coprod_\eta M_*(K_\eta) \arrow[uu] \arrow[dl] & & \mathcal{M}_* \arrow[ll] \arrow[uu] \arrow[dl] \\
\prod'_y M_*(\widehat{K}_y) \arrow[uu] & & \prod_y M_*(\widehat{\mathcal{O}}_y) \arrow[ll] \arrow[uu] &
\end{tikzcd}$$

where $\mathcal{M}_*$ *denotes the Zariski sheaf of the cycle module* M_*.

Proof. This is really just the definition of unramified cycles, see eq. 2.6. For $W^{02}_\bullet$ use Lemma 8. Note that certain finiteness conditions à la $\prod''$ and $\prod'''$ have disappeared here since they are automatically satisfied. □

Definition 3. *We define a complex of flasque sheaves (depending on the cycle module* M_* *which we have fixed):*

(4.5)
$$\begin{aligned}
\mathbf{I}_\bullet : 0 &\to \coprod_\eta M_*(K_\eta) \oplus \prod_y M_*(\widehat{\mathcal{O}}_y) \oplus \prod_x M_*(\widehat{\mathcal{O}}_x) \cdots \\
&\to \prod_y' M_*(\widehat{K}_y) \oplus \prod_x M_*(\widehat{\mathcal{O}}_x \otimes K) \oplus \prod_{x,y'_i} M_*(\widehat{\mathcal{O}}_{x,y'_i}) \cdots \\
&\to \prod_{x,y'_i}''' M_*(\widehat{K}_{x,y'_i}) \to 0.
\end{aligned}$$

(this is just the diagonal complex of fig. 4.4 with the initial term $\mathcal{M}_*$ *cut off as to make the second claim of the following proposition true)*

Proposition 5. *The diagonal complex of sheaves of the tricomplex* $H^0(W_\bullet)$, *fig. 4.4, is exact. In particular,*

$$0 \to \mathcal{M}_* \to \mathbf{I}_\bullet$$

is a flasque resolution of the sheaf $\mathcal{M}_*$.

For the proof of this we need some preparation:

Lemma 13. *The sequence of sheaves*

$$0 \longrightarrow \mathcal{M}_* \longrightarrow \coprod_\eta M_*(K_\eta) \oplus \prod_y M_*(\widehat{\mathcal{O}}_y) \longrightarrow \prod_y^{\prime+recip.} M_*(\widehat{K}_y) \longrightarrow 0$$

is exact.

This again shows that $\prod_y^{\prime+\text{recip.}} M_*(\widehat{K}_y)$ cannot be flasque, for otherwise the above complex would give a length 2 flasque resolution and so $H^2(X, \mathcal{M}_*) = 0$ always (which is not true).

Proof. The injectivity is clear since for each open set U the group of sections $\mathcal{M}_*(U)$ is defined as a subgroup of $\coprod_{\eta \in U^{(0)}} M_*(K_\eta)$, see eq. 2.6. In particular, injectivity holds on the level of stalks. Exactness in the middle is easy: Let x be an point in X, $U_x \ni x$ some open neighbourhood. Suppose local sections $\alpha_\eta \in M_*(K_\eta)$ and $\alpha_y \in M_*(\widehat{\mathcal{O}}_y)$ over U_x are given and go to zero on the right, i.e. for all codimension 1 points $y \in U_x^{(1)}$ we have $\operatorname{res}_{K_\eta}^{\widehat{K}_y} \alpha_\eta - \alpha_y = 0 \in M_*(\widehat{K}_y)$. Thus, taking the boundary at y in $\widehat{\mathcal{O}}_y$ we obtain

$$0 = \partial_y^{\widehat{K}_y} \operatorname{res}_{K_\eta}^{\widehat{K}_y} \alpha_\eta - \partial_y^{\widehat{K}_y} \alpha_y = \operatorname{res}_{\kappa(y)}^{\kappa(y)} \partial_y^K \alpha_\eta - \partial_y^{\widehat{K}_y} \alpha_y$$

by the compatibility of flat pullback (here already unwound to $\operatorname{res}_{K_\eta}^{\widehat{K}_y}$) with the differential in the Rost complex, axiom **R3a**. We also have $\partial_y^{\widehat{K}_y} \alpha_y = 0$ and so we deduce $\partial_y^K \alpha_\eta = 0$ since the restriction from $\kappa(y)$ to itself is the identity (to see this, use that $\operatorname{cor} \circ \operatorname{res}$ is multiplication with the extension degree, axiom **R2d**). However, this shows that $(\alpha_\eta)_\eta \in \mathcal{M}_*(U_x)$. Being a complex is obvious. For the surjectivity let x be a closed point in X and suppose we are given $(\alpha_y)_y$ in the right-hand side in some open neighbourhood U_x of x. For the (uncompleted) stalk $\mathcal{O}_x$ we have exactness of

$$0 \to M_*(\mathcal{O}_x) \to M_*(K_\eta) \to \coprod_{y \in (\mathcal{O}_x)^{(1)}} M_{*-1}(\kappa(y)) \to M_{*-2}(\kappa(x)) \to 0,$$

by Prop. 3. The element $(\partial_y^{\widehat{K}_y} \alpha_y)$ placed in the third term of this sequence (note that by the idelic condition $\prod'$ this is non-zero only for fin. many points y and the reciprocity constraint shows that the element goes to zero on the right) admits a preimage $f_\eta \in M_*(K_\eta)$. Secondly, note that (after possibly shrinking U_x to a smaller neighbourhood) $\beta_y := \operatorname{res}_{K_\eta}^{\widehat{K}_y} f_\eta - \alpha_y$ satisfies $\partial_y^{\widehat{K}_y} \beta_y = 0$ for all $y \in U_x$ by construction, so $\beta_y \in M_*(\widehat{\mathcal{O}}_y)$. It is easy to see that (f_η, β_y) is a preimage of α_y. This proves our claim. $\square$

Lemma 14. *The sequence of sheaves*

$$0 \to \prod_x M_*(\widehat{\mathcal{O}}_x) \to \prod_x M_*(\widehat{\mathcal{O}}_x \otimes K) \oplus \prod_{x, y_i'} M_*(\widehat{\mathcal{O}}_{x, y_i'}) \cdots$$
$$\cdots \to \prod_{x, y_i'}^{\prime\prime\prime + recip.} M_*(\widehat{K}_{x, y_i'}) \to 0$$

is exact.

Proof. The proof is very similar to the one of the previous lemma. For injectivity, Lemma 8 shows that the rows in the commutative diagram

(4.6)

$$\begin{tikzcd}
0 \arrow[r] & M_*(\widehat{\mathcal{O}}_x) \arrow[r] \arrow[d, phantom, "(\downarrow)"] & M_*(\widehat{K}_x) \arrow[r] \arrow[d, equal] & \coprod_{y' \in (\widehat{\mathcal{O}}_x)^{(1)}} M_{*-1}(\kappa(y')) \arrow[r] \arrow[d] & M_{*-2}(\kappa(x)) \\
0 \arrow[r] & M_*(\widehat{\mathcal{O}}_x \otimes K) \arrow[r] & M_*(\widehat{K}_x) \arrow[r] & \coprod_{\widehat{y}} M_{*-1}(\kappa(\widehat{y})) &
\end{tikzcd}$$

are exact, where $\widehat{y}$ runs through the transcendental primes in $\widehat{\mathcal{O}}_x$. The existence of the arrow ($\downarrow$) and its injectivity follow. For exactness in the middle, suppose we are given $\alpha_x \in M_*(\widehat{\mathcal{O}}_x \otimes K)$ and $\alpha_{x,y_i'} \in M_*(\widehat{\mathcal{O}}_{x,y_i'})$ going to zero on the right. This unwinds as

$$\forall x, y_i' : \operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_x} \alpha_x - \alpha_{x,y_i'} = 0 \in M_*(\widehat{K}_{x,y_i'}) \tag{4.7}$$

and applying the boundary $\partial^{\widehat{K}_{x,y_i'}}_{y_i'}$ yields $\operatorname{res}^{\kappa(y_i')}_{\kappa(y_i')} \partial^{\widehat{K}_x}_{y_i'} \alpha_x = \partial^{\widehat{K}_{x,y_i'}}_{y_i'} \alpha_{x,y_i'}$. We conclude $\partial^{\widehat{K}_x}_{y_i'} \alpha_x = 0$. By Lemma 7 the height 1 primes of $\widehat{\mathcal{O}}_x$ disjointly decompose into transcendental $\widehat{y}$ and non-transcendental y_i' primes. By $\alpha_x \in M_*(\widehat{\mathcal{O}}_x \otimes K)$ we thus have shown that the boundaries at *all* height 1 primes of $\widehat{\mathcal{O}}_x$ vanish and thus $\alpha_x \in M_*(\widehat{\mathcal{O}}_x)$ for each x. Finally, use eq. 4.7 to see that α_x maps to $\alpha_{x,y_i'}$ under the first arrow, showing that we have found the preimage. To see surjectivity, suppose we are given $\alpha_{x,y_i'} \in M_*(\widehat{K}_{x,y_i'})$. By the reciprocity constraint and the exactness of the first line of fig. 4.6 applied to $(\partial_{y_i'} \alpha_{x,y_i'}, 0_{\widehat{y}})_{y'}$ with y_i' running through the non-transcendental primes and 0 for the transcendental primes $\widehat{y}$, we obtain a preimage $f \in M_*(\widehat{K}_x)$. Since by construction $\partial_{\widehat{y}} f = 0$, the exactness of the lower row shows $f \in M_*(\widehat{\mathcal{O}}_x \otimes K)$. It is now obvious that $\beta_{x,y_i'} := \operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_x} f - \alpha_{x,y_i'} \in M_*(\widehat{\mathcal{O}}_{x,y_i'})$ and that $f \oplus \beta_{x,y_i'}$ provides a preimage. This completes the proof. $\square$

Proof (of Prop. 5). Combining Lemmata 13 and 14 we see that the diagonal complexes of the top and bottom face of the cube in fig. 4.4 (i.e. the opposite faces w.r.t. the arrows '$\uparrow$') are exact up to a cokernel isomorphic to $\coprod_x M_{*-2}(\kappa(x))$ because of the reciprocity constraints. It is not hard to see that this forces the overall diagonal complex to be exact. $\square$

Lemma 15. *The tricomplexes of sheaves $H^p(W_\bullet)$ for $p \geq 1$ are zero.*

Proof. Direct verification. $\square$

5. Comparison Maps for Idèles

Now we set up a bicomplex $\mathcal{B}$ from the *entries* of the tricomplexes $W_\bullet$,

- $W_\bullet$ being indexed by a lower index in $\{0, 1, 2\}$,
- the entries being indexed by an upper index in $\{0, 1, 2, 01, 02, 12, 012\}$ as in fig. 4.3:

$$\begin{tikzcd}
W_0^0 \oplus W_0^1 \oplus W_0^2 \arrow[r] \arrow[d] & W_1^0 \oplus W_1^1 \oplus W_1^2 \arrow[r] \arrow[d] & W_2^0 \oplus W_2^1 \oplus W_2^2 \arrow[r] \arrow[d] & 0 \\
W_0^{01} \oplus W_0^{02} \oplus W_0^{12} \arrow[r] \arrow[d] & W_1^{01} \oplus W_1^{02} \oplus W_1^{12} \arrow[r] \arrow[d] & W_2^{01} \oplus W_2^{02} \oplus W_2^{12} \arrow[r] \arrow[d] & 0 \\
W_0^{012} \arrow[r] & 0 \arrow[r] & 0 \arrow[r] & 0,
\end{tikzcd}$$

i.e. the columns are the diagonal complexes of the tricomplexes $W_\bullet$, truncated from the top by cutting away the initial term $W_\bullet^X$ ('un-augmented'); and the rows follow $W_0 \to W_1 \to W_2$. Clearly $\mathcal{B}$ is a bicomplex of sheaves with values in abelian groups. Apply the global section functor $\Gamma(X, \cdot)$ and note that this preserves exactness since all sheaves in the bicomplex are flasque.

We quickly study the two bicomplex spectral sequences associated to $\Gamma(X, \mathcal{B})$. We denote by ${}^{\downarrow}E^{\bullet}$ the spectral sequence whose E^0-page differential comes from the downward arrows '↓', and by ${}^{\rightarrow}E^{\bullet}$ the spectral sequence whose E^0-page differential comes from '→'.

Proposition 6 (Gersten-to-Idèles bicomplex)**.** *The bicomplex $\Gamma(X, \mathcal{B})$ has*

$$\begin{aligned} {}^{\downarrow}E^2_{p,0} &= H^p(\Gamma(X, \mathbf{I}_\bullet)) \qquad \textit{and} \qquad {}^{\downarrow}E^2_{p,q} = 0 \textit{ for } q \geq 1 \\ {}^{\rightarrow}E^2_{0,q} &= A^q(X, M_*) \qquad \textit{and} \qquad {}^{\rightarrow}E^2_{p,q} = 0 \textit{ for } p \geq 1 \end{aligned}$$

and these pages are stationary (= E^∞ respectively). In particular, it trivially converges to $H^p(X, \mathcal{M}_*)$.

Here the groups $A^q(X, M_*)$ are as in [**26**, §5, especially p. 356; first paragraph].

Proof. For ${}^{\downarrow}E^1$ we find in the first row

$$0 \to \Gamma(X, W_0^X) \to \Gamma(X, W_1^X) \to \Gamma(X, W_2^X) \to 0$$

and zero elsewhere. This follows from the observation that the diagonal complex each tricomplex W_i is exact as sheaves, and then also after applying $\Gamma(X, \cdot)$ by flasqueness. Unwinding definitions this yields Rost's Gersten resolution of $\mathcal{M}_*$, namely

$$0 \to \coprod_{\eta \in X^{(0)}} M_*(K_\eta) \overset{\partial^\eta_y}{\to} \coprod_{y \in X^{(1)}} M_{*-1}(\kappa(y)) \overset{\partial^y_x}{\to} \coprod_{x \in X^{(2)}} M_{*-2}(\kappa(x)) \to 0,$$

i.e. in the ${}^{\downarrow}E^2 = {}^{\downarrow}E^\infty$-page the first row are the corresponding cohomology groups, denoted by $A^q(X, M_*)$. Contrastingly, the ${}^{\rightarrow}E^1$-page reads

$$\begin{tikzcd} \coprod_{\eta \in X^{(0)}} M_*(K_\eta) \oplus \prod_{y \in X^{(1)}} M_*(\widehat{\mathcal{O}}_y) \oplus \prod_{x \in X^{(2)}} M_*(\widehat{\mathcal{O}}_x) \arrow[r] \arrow[d] & 0 \arrow[d] & \cdots \\ \prod'_{y \in X^{(1)}} M_*(\widehat{K}_y) \oplus \prod_{x \in X^{(2)}} M_*(\widehat{\mathcal{O}}_x \otimes K) \oplus \prod_{x \in X^{(2)}, y'_i} M_*(\widehat{\mathcal{O}}_{x,y'_i}) \arrow[r] \arrow[d] & 0 \arrow[d] & \cdots \\ \prod'''_{x \in X^{(2)}, y'_i} M_*(\widehat{K}_{x,y'_i}) \arrow[r] & 0 & \cdots \end{tikzcd}$$

which just uses Prop. 5 in the left-most column and Lemma 15 in the other two. Again use that $\Gamma(X, \cdot)$ preserves exactness on flasque sheaves. For each x the primes y'_i run through the non-transcendental height 1 primes of $\widehat{\mathcal{O}}_x$. It is obvious that the left-most column agrees with $\mathbf{I}_\bullet$ as given in eq. 4.5. □

Based on this result, we can formulate explicit comparison maps between the groups $A^q(X, M_*)$ and $H^q(\Gamma(X, \mathbf{I}_\bullet))$:

Proposition 7. *The comparison isomorphism*

$$\varphi : A^1(X, M_*) \to H^1(\Gamma(X, \mathbf{I}_\bullet))$$

is given by sending a cycle $\sum [y, \gamma_y] \in C^1(X, M_*)$ with $y \in X^{(1)}$ and $\gamma_y \in M_{*-1}(\kappa(y))$ to the idèle

$$\begin{aligned} (\alpha_y, \alpha_x, \alpha_{x,y'_i}) \in {\prod}'_{y \in X^{(1)}} M_*(\widehat{K}_y) &\oplus \prod_{x \in X^{(2)}} M_*(\widehat{\mathcal{O}}_x \otimes K) \\ &\oplus \prod_{x \in X^{(2)}, y'_i} M_*(\widehat{\mathcal{O}}_{x,y'_i}) \end{aligned}$$

$$\alpha_y = f_y, \qquad \alpha_x = g_x, \qquad \alpha_{x,y'_i} = \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_x} g_x - \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} f_y,$$

where

- *f_y is any preimage of γ_y under $M_*(\widehat{K}_y) \to M_{*-1}(\kappa(y))$,*
- *g_x is any preimage of $(\operatorname{res}_{\kappa(y)}^{\kappa(y')} \gamma_y)_y$ under $M_*(\widehat{K}_x) \to \coprod_{y'} M_{*-1}(\kappa(y'))$.*

In particular, these preimages exist. This defines φ on generators and does not depend on any choices (all idèles produced this way are cohomologous in $\mathbf{I}_\bullet$).

Proof. This is a standard argument proceeding by unwinding zig-zags in the bicomplex. We follow the zig-zag

$$\begin{tikzcd} W_0^0 \oplus W_0^1 \oplus W_0^2 \arrow[r] \arrow[d] & W_1^0 \oplus W_1^1 \oplus W_1^2 \\ W_0^{01} \oplus W_0^{02} \oplus W_0^{12}. & \end{tikzcd} \tag{5.1}$$

In our case, we begin with $\sum [y, \gamma_y] \in \coprod_y M_{*-1}(\kappa(y)) = C^1(X, M_*) = W_1^X$. In the sense of an edge map, this kernel maps diagonally to an element in the bicomplex $\mathcal{B}$, $W_1^X \to W_1^0 \oplus W_1^1 \oplus W_1^2$ in the upper-right of our zig-zag, namely

$$[y, \gamma_y] \in \coprod\nolimits_y M_{*-1}(\kappa(y)) \qquad \text{and} \qquad [y, \operatorname{res}_{\kappa(y)}^{\kappa(y')} \gamma_y] \in \prod\nolimits_x'' \coprod\nolimits_{y'} M_{*-1}(\kappa(y'))$$

in $W_1^1 \oplus W_1^2$ (note that $W_1^0 = 0$). As is easily confirmed, a preimage in $W_0^0 \oplus W_0^1 \oplus W_0^2$ is given by

$$0 \in \coprod\nolimits_\eta M_*(K_\eta), \qquad [\widehat{K}_y, f_y] \in \prod\nolimits_y' M_*(\widehat{K}_y), \qquad [\widehat{K}_x, g_x] \in \prod\nolimits_x'' M_*(\widehat{K}_x)$$

with f_y and g_x as stated in the claim. The existence of f_y follows from the exactness of $M_*(\widehat{K}_y) \to M_{*-1}(\kappa(y)) \to 0$ (Gersten for $\widehat{\mathcal{O}}_y$, Prop. 4). For the existence of g_x one uses the exactness of

$$M_*(K_\eta) \longrightarrow \coprod_y M_{*-1}(\kappa(y)) \longrightarrow \coprod_x M_{*-2}(\kappa(x)) \tag{5.2}$$

(Gersten for $\mathcal{O}_x$, Prop. 3), noting that $\sum [y, \gamma_y]$ goes to zero on the right which holds by the assumption it represents an element in $A^1(X, M_*)$, i.e. it is a cocycle. We get a preimage $\tilde{g} \in M_*(K_\eta)$ and then define $g_x := \operatorname{res}_{K_\eta}^{\widehat{K}_x} \tilde{g}$. It is easy to see that this element has the desired properties. Under the downward differential in fig. 5.1, to $W_0^{01} \oplus W_0^{02} \oplus W_0^{12}$, we find

$$\alpha_y = f_y, \qquad \alpha_x = g_x, \qquad \alpha_{x,y_i'} = \operatorname{res}_{\widehat{K}_x}^{\widehat{K}_{x,y_i'}} g_x - \operatorname{res}_{\widehat{K}_y}^{\widehat{K}_{x,y_i'}} f_y$$

and it is easy to see that the differential $W_0^{01} \oplus W_0^{02} \oplus W_0^{12} \to W_0^{012}$ sends this element to zero, so this gives us a representative for $H^1(\Gamma(X, \mathbf{I}_\bullet))$. It follows from extensive diagram chasing that this definition is well-defined, a homomorphism, and an isomorphism. □

Proposition 8. *The comparison isomorphism*

$$\theta : H^2(\Gamma(X, \mathbf{I}_\bullet)) \to A^2(X, M_*)$$

is given by sending $(\alpha_{x,y_i'})_{x,y_i'}$ with $\alpha_{x,y_i'} \in \prod'''_{x.y_i'} M_*(\widehat{K}_{x,y_i'})$ to

$$\left(\sum\nolimits_{y_i'} \partial_x^{y_i'} \partial_{y_i'} \alpha_{x,y_i'} \right)_x \in \coprod\nolimits_{x \in X^{(2)}} M_{*-2}(\kappa(x)),$$

where for every $x \in X^{(2)}$ the parameter y_i' runs through all non-transcendental primes in $\widehat{\mathcal{O}}_x$.

PROOF. As in the previous proof, this is an immediate consequence of chasing zig-zags in the bicomplex. This time the relevant zig-zag is

$$\begin{tikzcd}
 & W_1^0 \oplus W_1^1 \oplus W_1^2 \arrow[r] \arrow[d] & W_2^0 \oplus W_2^1 \oplus W_2^2 \\
W_0^{01} \oplus W_0^{02} \oplus W_0^{12} \arrow[r] \arrow[d] & W_1^{01} \oplus W_1^{02} \oplus W_1^{12} & \\
W_0^{012}. & &
\end{tikzcd}$$

□

6. Comparison Maps for Čech cohomology

Firstly, recall that for an open cover $\mathfrak{U} = (U_\alpha)_{\alpha \in I}$ and Zariski sheaf $\mathcal{F}$ there are Čech cohomology groups, which we denote by $\check{H}^i(\mathfrak{U}, \mathcal{F})$, defined as the cohomology of the Čech complex

$$\check{\mathrm{C}}^i(\mathfrak{U}, \mathcal{F}) := \prod_{\alpha_0 \dots \alpha_r \in I^{i+1}} \mathcal{F}(U_{\alpha_0 \dots \alpha_r}), \qquad \delta : \check{\mathrm{C}}^i(\mathfrak{U}, \mathcal{F}) \to \check{\mathrm{C}}^{i+1}(\mathfrak{U}, \mathcal{F}),$$

where we denote by U_α an open in $\mathfrak{U}$, and by $U_{\alpha_0 \dots \alpha_r} := \bigcap_{i=0,\dots,r} U_{\alpha_i}$ the respective intersections. For any refinement $\mathfrak{U}'$ of $\mathfrak{U}$, there is a canonical induced morphism $\check{H}^i(\mathfrak{U}, \mathcal{F}) \to \check{H}^i(\mathfrak{U}', \mathcal{F})$ (and 'unique' on $\check{\mathrm{C}}^\bullet$ up to homotopies). Finally, $\check{H}^i(X, \mathcal{F})$ is defined as $\operatorname{colim}_{\mathfrak{U}} \check{H}^i(X, \mathcal{F})$ over the diagram in which $\mathfrak{U}$ runs through all open covers[5] and arrows are the refinements. This is a filtering colimit since any two covers admit a common refinement. We have $\check{H}^i(X, \mathcal{F}) \cong H^i(X, \mathcal{F})$ for all sheaves we work with (as they all admit a flasque resolution), so we may express sheaf cohomology this way.

A cocycle $(f_{\alpha_0 \dots \alpha_i})_{\alpha_0 \dots \alpha_i} \in \check{\mathrm{C}}^i(\mathfrak{U}, \mathcal{F})$ is called *alternating* if $f_{\sigma(\alpha_0) \dots \sigma(\alpha_i)} = (-1)^{\operatorname{sgn} \sigma} f_{\alpha_0 \dots \alpha_i}$ holds for all permutations $\sigma \in S_{i+1}$ and $f_{\alpha_0 \dots \alpha_w \alpha_w \dots \alpha_{i-1}} = 0$ for all $0 \leq w \leq i-1$. The alternating Čech cocycles form a subcomplex of $\check{\mathrm{C}}^i(\mathfrak{U}, \mathcal{F})$ and one can show that the inclusion of the subcomplex is a quasi-isomorphism.

Remark 3. *It is worth emphasizing that* $f_{\alpha_0 \dots \alpha_w \alpha_w \dots \alpha_{i-1}} = 0$ *is a separate condition. It follows from the first property just for elements which are not* 2*-torsion, à la* $f_{\alpha\beta} = -f_{\beta\alpha}$*, so* $2 f_{\alpha\alpha} = 0$ *in the case of* $i = 1$*.*

Remark 4. *The explicit determination of Čech cohomology representatives for sections of the sheaf* $\mathcal{K}_2$ *has served as a key tool in* [**4**, Prop. 2.8].

In this text we shall always work with this subcomplex of alternating cochains.

Lemma 16 (Algebraic "Partition of Unity")**.** *Assume the open cover* $\mathfrak{U}$ *is finite, i.e.* I *is a finite set. Suppose* $\mathcal{F}$ *is a flasque sheaf such that*

(1) *for every restriction* $\operatorname{res}^U_V : \mathcal{F}(U) \to \mathcal{F}(V)$ *for any two opens of the form* $U_{\gamma_0 \dots \gamma_s \gamma_{s+1} \dots \gamma_r} =: V \subseteq U := U_{\gamma_0 \dots \gamma_s}$ *there is a morphism*

$$E^U_V : \mathcal{F}(V) \to \mathcal{F}(U)$$

(2) *for every open* $V := U_{\gamma_0 \dots \gamma_s}$ *we have*

$$\sum_{\alpha \in I} E^V_{V_\alpha} \circ \operatorname{res}^V_{V_\alpha} = \mathrm{id}_{\mathcal{F}(V)}, \qquad \text{where } V_\alpha := V \cap U_\alpha$$

---

[5] We suppress some set-theoretical issues.

(3) *for every open* $V := U_{\gamma_0\dots\gamma_s}$ *and all indices* $\beta \in I$ *we have*

$$\coprod_{\alpha\in I}\mathcal{F}(V_\alpha)\to\mathcal{F}(V_\beta),\qquad \textit{where } V_\alpha := V\cap U_\alpha .$$

$$\sum_{\alpha\in I}\operatorname{res}^{V}_{V_\beta}\circ E^{V}_{V_\alpha} = \sum_{\alpha\in I} E^{V_\beta}_{V_{\alpha\beta}}\circ\operatorname{res}^{V_\alpha}_{V_{\alpha\beta}} .$$

Define a homomorphism

$$H : \check{\mathrm{C}}^{i}\left(\mathfrak{U},\mathcal{F}\right)\to\check{\mathrm{C}}^{i-1}\left(\mathfrak{U},\mathcal{F}\right)$$

$$(Hf)_{\beta_0\dots\beta_{i-1}} := \sum_{\alpha\in I} E^{U_{\beta_0\dots\beta_{i-1}}}_{U_{\alpha\beta_0\dots\beta_{i-1}}} f_{\alpha\beta_0\dots\beta_{i-1}} .$$

Then H *is a contracting homotopy for the Čech complex* $\check{\mathrm{C}}^{\bullet}\left(\mathfrak{U},\mathcal{F}\right)$*, i.e.*

$$H\delta+\delta H = \mathrm{id}_{\check{\mathrm{C}}^{i}(\mathfrak{U},\mathcal{F})} .$$

Proof. Direct computation. □

Lemma 17. *Assume the open cover* $\mathfrak{U}$ *is finite, i.e.* I *is a finite set. For all the sheaves as in eq. 4.1 the assumptions of Lemma 16 can be fulfilled: For* $\prod_y A_y$ *with* A_y *some abelian group (e.g.* $M_*(\widehat{K}_y)$*,* $M_*(\kappa(y))$*, etc.) fix a disjoint decomposition*

$$X^{(1)} = \dot{\bigcup_{\alpha\in I}}\Sigma_\alpha \textit{ with } \Sigma_\alpha \subseteq U^{(1)}_\alpha . \tag{6.1}$$

Define for any open $U := U_{\gamma_0\dots\gamma_s}$ *and further intersection* $V := U\cap U_\beta$ *(*$= U_{\gamma_0\dots\gamma_s\beta}$*) the homomorphism*

$$E^U_V : \mathcal{F}(V)\to\mathcal{F}(U)$$

$$\prod_{y\in V^{(1)}} A_y \mapsto \prod_{y\in U^{(1)}} A_y \qquad \textit{by} \qquad \begin{cases} \mathrm{id}_{A_y} & \textit{if } y\in\Sigma_\beta \\ 0 & \textit{otherwise.}\end{cases}$$

and transitively define $E^U_V := E^U_{U\cap U_\beta}\circ E^{U\cap U_\beta}_V$ *if* $V := U\cap U_\beta\cap U_{\beta_2}$ *etc. (Use* $X^{(0)}$ *and* $U^{(0)}_\alpha$*, or* $X^{(2)}$ *and* $U^{(2)}_\alpha$ *for sheaves of the types* $\coprod_\eta A_\eta$ *or* $\prod_x A_x$ *respectively. For* $\prod_{x,y'_i}$ *also use* $X^{(2)}$ *and* $U^{(2)}_\alpha$ *and recall that for every* x *the parameter* y' *just runs through certain primes of* $\widehat{\mathcal{O}}_x$*).*

Proof. A disjoint decomposition as in eq. 6.1 exists since $\mathfrak{U}$ is an open cover, so $X = \bigcup U_\alpha$; in particular $X^{(i)} = \bigcup U^{(i)}_\alpha$. For example (well-)order the set I and then let $x\in\Sigma_\alpha$ if and only if α is the (unique) smallest element of I such that $x\in U_\alpha$. It remains to prove the properties (2) and (3). For (2) observe that for every prime y the map $E^V_{V_\alpha}\circ\operatorname{res}^V_{V_\alpha}$ on A_y is id_{A_y} if $y\in\Sigma_\alpha$ and zero otherwise. Since $\mathfrak{U}$ is a finite cover and the Σ_α form a *disjoint* decomposition, the equality

$$\sum_{\alpha\in I} E^V_{V_\alpha}\circ\operatorname{res}^V_{V_\alpha} = \mathrm{id}_{\mathcal{F}(V)}$$

follows, proving property (2). Then (3) is a direct verification. □

We now use these contracting homotopies to work out certain *very explicit* maps between various cohomology groups:

Proposition 9. *Let X be integral and η denote its generic point. Suppose $\mathfrak{U} = (U_\alpha)_{\alpha\in I}$ is a finite open cover of X and fix a disjoint decomposition $X = \dot{\bigcup}_{\alpha\in I} \Sigma_\alpha$ as in Lemma 17. Then the canonical morphism*[6]

$$\tau : \check{H}^1(\mathfrak{U}, \mathcal{M}_*) \to H^1(\Gamma(X, \mathbf{I}_\bullet))$$

admits the following explicit description: If $f := (f_{\alpha\beta})_{\alpha\beta} \in \check{\mathrm{C}}^1(\mathfrak{U}, \mathcal{M}_*)$ is a Čech 1-cocycle, define an idèle $\tilde{f}$ by

$$\tilde{f}^{01}_y := -\operatorname{res}^{\widehat{K}_y}_{K_\eta} f_{\beta_\eta\beta_y} \qquad \tilde{f}^{02}_x := -\operatorname{res}^{\widehat{K}_x}_{K_\eta} f_{\beta_\eta\beta_x} \qquad \tilde{f}^{12}_{x,y'_i} := -\operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} f_{\beta_y\beta_x},$$

where $\beta_\eta, \beta_y, \beta_x \in I$ are the unique indices such that $\eta \in \Sigma_{\beta_\eta}$, $y \in \Sigma_{\beta_y}$, $x \in \Sigma_{\beta_x}$ and y in the right-most formula is the image of y'_i under $\operatorname{Spec}\widehat{\mathcal{O}}_x \to \operatorname{Spec}\mathcal{O}_x$. Then $[\tau f] = [\tilde{f}]$ as cohomology classes.

Proof. We build a bicomplex whose p-th row is built by taking the p-th Čech resolution group of a copy of $\mathbf{I}_\bullet$. Again we follow zig-zags as in the proof of Prop. 7, but now using a bicomplex comparing the Čech and the idèle resolution. The groups in

$$\begin{tikzcd}
 & & & {}^{\downarrow}E^2_{1,0} \\
 & & & {}^{\downarrow}E^1_{1,0} \arrow[u, Rightarrow] \\
 & & E^0_{0,0} \arrow[r] \arrow[d] & E^0_{1,0} \arrow[u, Rightarrow] \\
{}^{\rightarrow}E^2_{0,1} \arrow[r, Rightarrow] & {}^{\rightarrow}E^1_{0,1} \arrow[r, Rightarrow] & E^0_{0,1} &
\end{tikzcd}$$

are just

$$\begin{tikzcd}
 & & & H^1(\Gamma(X, \mathbf{I}_\bullet)) \\
 & & & \Gamma(X, \mathbf{I}_1) \arrow[u, Rightarrow] \\
 & & \check{\mathrm{C}}^0(\mathfrak{U}, \mathbf{I}_0) \arrow[r] \arrow[d] & \check{\mathrm{C}}^0(\mathfrak{U}, \mathbf{I}_1) \arrow[u, Rightarrow] \\
\check{H}^1(\mathfrak{U}, \mathcal{M}_*) \arrow[r, Rightarrow] & \check{\mathrm{C}}^1(\mathfrak{U}, \mathcal{M}_*) \arrow[r, Rightarrow] & \check{\mathrm{C}}^1(\mathfrak{U}, \mathbf{I}_0). &
\end{tikzcd}$$

Explicitly, we choose any representative for our Čech cohomology class $(f_{\alpha\beta})_{\alpha\beta} \in \check{\mathrm{C}}^1(\mathfrak{U}, \mathcal{M}_*)$ in ${}^{\rightarrow}E^1_{0,1}$. We use the edge map to get to $E^0_{0,1}$. We construct a preimage in $E^0_{0,0}$ by using the contracting homotopy of the Čech complex of Lemma 16. This yields an element in $E^0_{0,0}$

$$f'_\beta := \sum_{\alpha\in I} E^{U_\beta}_{U_{\alpha\beta}} f_{\alpha\beta} \text{ with } (f'_\beta) \in \check{\mathrm{C}}^1(\mathfrak{U}, \coprod_\eta M_*(K_\eta) \oplus \coprod_y M_*(\widehat{\mathcal{O}}_y) \oplus \coprod_x M_*(\widehat{\mathcal{O}}_x)).$$

Explicitly,

$$f'^0_\beta := \sum_\alpha \delta_{\eta\in\Sigma_\alpha} f_{\alpha\beta} \qquad f'^1_\beta := \sum_\alpha \delta_{y\in\Sigma_\alpha} \operatorname{res}^{\widehat{K}_y}_{K_\eta} f_{\alpha\beta} \qquad f'^2_\beta := \sum_\alpha \delta_{x\in\Sigma_\alpha} \operatorname{res}^{\widehat{K}_x}_{K_\eta} f_{\alpha\beta}$$

---

[6] isomorphism if we replace the left-hand side by the colimit over all refinements

where $\delta_A = \mathrm{id}$ if A is a true expression, 0 otherwise. Under the differential $E^0_{0,0} \to E^0_{1,0}$ this maps to the element $(f''_\beta)_\beta \in E^0_{1,0} = \check{\mathrm{C}}^0(\mathfrak{U}, \mathbf{I}_1)$ given by

$$f''^{01}_\beta := f'^1_\beta - \operatorname{res}^{\widehat{K}_y}_{K_\eta} f'^0_\beta = \sum_\alpha (\delta_{y\in\Sigma_\alpha} - \delta_{\eta\in\Sigma_\alpha}) \operatorname{res}^{\widehat{K}_y}_{K_\eta} f_{\alpha\beta}$$

$$f''^{02}_\beta := f'^2_\beta - \operatorname{res}^{\widehat{K}_x}_{K_\eta} f'^0_\beta = \sum_\alpha (\delta_{x\in\Sigma_\alpha} - \delta_{\eta\in\Sigma_\alpha}) \operatorname{res}^{\widehat{K}_x}_{K_\eta} f_{\alpha\beta}$$

$$f''^{12}_\beta := \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_x} f'^2_\beta - \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} f'^1_\beta = \sum_\alpha (\delta_{x\in\Sigma_\alpha} - \delta_{y\in\Sigma_\alpha}) \operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} f_{\alpha\beta},$$

where y is the image of y'_i under $\operatorname{Spec} \widehat{\mathcal{O}}_x \to \operatorname{Spec} \mathcal{O}_x$. This is a Čech 0-cocycle in $\check{H}^0(\mathfrak{U}, \mathbf{I}_1)$, so it glues to a global section in $\Gamma(X, \mathbf{I}_1)$. Thus, if we want to evaluate the component of f''^{01}_β at a point $y \in X^{(1)}$, i.e. the element $f''^{01}_{\beta,y}$ in

$$\Gamma(U_\beta, W^{01}_1) \ni f'^{01}_\beta = (f'^{01}_{\beta,y})_y \in {\prod}'_{y\in U^{(1)}_\beta} M_*(\widehat{K}_y),$$

we need to choose for β any index in I such that $y \in U^{(1)}_\beta$ (and because it glues to a global section, the choice of any such β does not matter). Similarly for the components $f''^{02}_{\beta',x}$, f''^{12}_{β',x,y'_i}, we have to pick any index $\beta' \in I$ such that $x \in U^{(2)}_{\beta'}$ (note that in the case of f''^{12}_{β',x,y'_i} the argument y'_i runs through non-transcendental height 1 primes of $\widehat{\mathcal{O}}_x$ independently of any open on which we evaluate the sheaf, so only x matters for the choice of the index β'). As we had already picked a disjoint decomposition $\dot{\bigcup} \Sigma_\alpha = X$, we may also use it to find canonical indices β, β' for these purposes. For any $y \in X^{(1)}$ let $\beta_y \in I$ be the unique index such that $y \in \Sigma_{\beta_y}$. We get

$$f''^{01}_{\beta_y,y} = \sum_\alpha (\delta_{y\in\Sigma_\alpha} - \delta_{\eta\in\Sigma_\alpha}) \operatorname{res}^{\widehat{K}_y}_{K_\eta} f_{\alpha\beta_y} = \operatorname{res}^{\widehat{K}_y}_{K_\eta} f_{\beta_y\beta_y} - \operatorname{res}^{\widehat{K}_y}_{K_\eta} f_{\beta_\eta\beta_y}$$

since $\delta_{y\in\Sigma_\alpha}$ will only be non-zero for $\alpha = \beta_y$ by uniqueness and there is just a single β_η with $\eta \in \Sigma_\eta$ (since we assume X is integral, so it is irreducible with a unique generic point $X^{(0)} = \{\eta\}$). Finally, as we work with alternating Čech cochains, we must have $f_{\beta_y\beta_y} = 0$ - see Rmk. 3 for 2-torsion. For W^{02}_1 and W^{12}_1 one proceeds analogously, giving our claim. By homological generalities the analogous map is an isomorphism if we replace $\check{H}^1(\mathfrak{U}, \mathcal{M}_*)$ by the colimit over all refinements, i.e. $H^1(X, \mathcal{M}_*)$. □

Proposition 10. *Let X be integral and η denote its generic point. Suppose $\mathfrak{U} = (U_\alpha)_{\alpha\in I}$ is a finite open cover of X and fix a disjoint decomposition $X = \dot{\bigcup}_{\alpha\in I} \Sigma_\alpha$ as in Lemma 17. The canonical morphism*[7]

$$\lambda : \check{H}^2(\mathfrak{U}, \mathcal{M}_*) \to A^2(X, M_*)$$

can be described as follows: For $f := (f_{\alpha\beta\gamma})_{\alpha\beta\gamma} \in \check{\mathrm{C}}^2(\mathfrak{U}, \mathcal{M}_*)$ define

$$h_x := \sum_{x\in\overline{\{y\}}} \partial^y_x \partial^\eta_y f_{\beta_\eta\beta_y\beta_x} \in M_{*-2}(\kappa(x)), \tag{6.2}$$

where

---

[7] isomorphism if we replace the left-hand side by the colimit over all refinements

- *β_x is to be chosen such that $x \in U_{\beta_x}$,*
- *$\beta_y \in I$ is the unique index such that $y \in \Sigma_{\beta_y}$, β_η the unique element such that $\eta \in \Sigma_{\beta_\eta}$.*

Despite true in this more general version, the statement of the lemma becomes entirely symmetric if we also choose $\beta_x \in I$ as the unique element with $x \in \Sigma_{\beta_x}$.

Proof. This follows from a diagram chase in the bicomplex whose columns come from the Čech resolution applied to rows built from Rost's Gersten resolution. Then the two E^1-pages, ${}^{\downarrow}E^1$ and ${}^{\rightarrow}E^1$, give the original Rost Gersten resolution in the top row and zero elsewhere, and the Čech resolution of $\mathcal{M}_*$ in the left column respectively. Both spectral sequences degenerate at the E^2-page and a diagram chase along a zig-zag provides us with an explicit description of the isomorphism in question. We abbreviate $\check{C}^2(\mathcal{M}_*) := \check{C}^2(\mathfrak{U}, \mathcal{M}_*)$. More precisely, we run along the zig-zag

$$\begin{tikzcd}
 & & & & {}^{\downarrow}E^2_{2,0} & = A^2(X, M_*) \\
 & & & & {}^{\downarrow}E^1_{2,0} \arrow[u, Rightarrow] & = \coprod M_{*-2}(\kappa(x)) \\
 & & & E^0_{1,0} \arrow[r] \arrow[d] & E^0_{2,0} \arrow[u, Rightarrow] & \\
\check{H}^2(\mathcal{M}_*) & \check{C}^2(\mathcal{M}_*) & E^0_{0,1} \arrow[r] \arrow[d] & E^0_{1,1} & & \\
{}^{\rightarrow}E^2_{0,2} \arrow[u, equal] \arrow[r, Rightarrow] & {}^{\rightarrow}E^1_{0,2} \arrow[u, equal] \arrow[r, Rightarrow] & E^0_{0,2} & & &
\end{tikzcd}$$

□

7. Products

One can consider pairings of cycle modules, as defined in [**26**, Def. 2.1]. Such a pairing consists of a bilinear pairing of abelian groups

$$\cdot : M_p(F) \times M'_q(F) \longrightarrow M''_{p+q}(F),$$

where M, M', M'' are cycle modules. Certain axioms, called **P1-P3** in [**26**], need to be fulfilled. We are only interested in the case $M = M' = M''$, i.e. a 'product' operation

$$\cdot : M_p(F) \times M_q(F) \longrightarrow M_{p+q}(F). \tag{7.1}$$

Example 6. *For example $M_* = K^M_*$ has such a pairing, just given by the ordinary product in the Milnor K theory ring.*

7.1. Cup Product on Čech Cochains.

We quickly recall the construction of the cup product in sheaf cohomology on the Čech resolution. For general sheaves $\mathcal{F}, \mathcal{G}$ with values in abelian groups, the tensor sheaf $\mathcal{F} \otimes_{\underline{\mathbf{Z}}} \mathcal{G}$ (where $\underline{\mathbf{Z}}$ denotes the locally constant sheaf $\mathbf{Z}$) has stalks

$$(\mathcal{F} \otimes_{\underline{\mathbf{Z}}} \mathcal{G})_x = \mathcal{F}_x \otimes_{\underline{\mathbf{Z}}_x} \mathcal{G}_x = \mathcal{F}_x \otimes_{\mathbf{Z}} \mathcal{G}_x.$$

For Čech cochains on an open cover $\mathfrak{U} = (U_\alpha)_{\alpha \in I}$ ones fixes an ordering on the index set I and defines a $\mathbf{Z}$-bilinear pairing

$$\smile : \check{C}^p(\mathfrak{U}, \mathcal{F}) \times \check{C}^q(\mathfrak{U}, \mathcal{G}) \to \check{C}^{p+q}(\mathfrak{U}, \mathcal{F} \otimes_{\underline{\mathbf{Z}}} \mathcal{G}) \tag{7.2}$$

$$(f \smile g)_{\alpha_0 \ldots \alpha_{p+q}} := {}^{\mathcal{F}}\mathrm{res}^{U_{\alpha_0 \cdots \alpha_p}}_{U_{\alpha_0 \cdots \alpha_{p+q}}} f_{\alpha_0 \ldots \alpha_p} \otimes {}^{\mathcal{G}}\mathrm{res}^{U_{\alpha_p \cdots \alpha_{p+q}}}_{U_{\alpha_0 \cdots \alpha_{p+q}}} g_{\alpha_p \ldots \alpha_{p+q}},$$

where $\alpha_0 < \cdots < \alpha_{p+q}$ and where ${}^{\mathcal{F}}\text{res}$ and ${}^{\mathcal{G}}\text{res}$ denote the restrictions to smaller opens of the sheaves $\mathcal{F}$, $\mathcal{G}$ respectively; and are not to be confused with the restrictions of a cycle module. Then prolong $(f \smile g)_{\alpha_0 \dots \alpha_{p+q}}$ to an alternating Čech cochain by sign-permuting indices to $\alpha_0 < \cdots < \alpha_{p+q}$.[8] The identity

$$\delta(f \smile g) = \delta f \smile g + (-1)^p \, f \smile \delta g$$

is easy to show and proves that eq. 7.2 induces a pairing of Čech cohomology groups, the *cup product.*

Remark 5. *If one defines the cup product in a derived setting as the morphism ‘$\cup$’ in*

$$\smile: \mathbf{R}\Gamma(X, \mathcal{F}) \otimes^{\mathbf{L}}_{\mathbf{Z}} \mathbf{R}\Gamma(X, \mathcal{G}) \overset{\cup}{\to} \mathbf{R}\Gamma(X, \mathcal{F} \otimes^{\mathbf{L}}_{\underline{\mathbf{Z}}} \mathcal{G}) \overset{\mathbf{R}\Gamma(pr)}{\to} \mathbf{R}\Gamma(X, \mathcal{F} \otimes_{\underline{\mathbf{Z}}} \mathcal{G}),$$

this cup product relates (after taking the colimit over all refinements of covers) to the one in eq. 7.2 by composing with $\mathbf{R}\Gamma(pr)$, where $pr : \mathcal{F} \otimes^{\mathbf{L}}_{\underline{\mathbf{Z}}} \mathcal{G} \to \mathcal{F} \otimes_{\underline{\mathbf{Z}}} \mathcal{G}$ is the natural morphism.

Next, fix a cycle module M_* along with a pairing $\cdot : M_n \times M_m \to M_{n+m}$ to itself as in the previous subsection. This induces a $\underline{\mathbf{Z}}$-bilinear pairing

$$\begin{aligned} \cdot : \mathcal{M}_n \otimes_{\underline{\mathbf{Z}}} \mathcal{M}_m &\longrightarrow \mathcal{M}_{n+m} \\ e \otimes f &\longmapsto e \cdot f \qquad \text{(the pairing of cycle modules)} \end{aligned}$$

of $\underline{\mathbf{Z}}$-module sheaves, i.e. sheaves with values in abelian groups. Recall that for every open $U \subseteq X$ the group $\mathcal{M}_*(U)$ is defined as in eq. 2.6. Thus, showing that the pairing is well-defined amounts to proving that $e \cdot f$ is unramified if e, f are, but by axiom **P3** we have for every discrete valuation v the identity

$$\partial_v(e \cdot f) = \partial_v(e) \cdot s_v^{\pi}(f) + (-1)^n \, s_v^{\pi}(e) \cdot \partial_v(f) + \{-1\} \cdot \partial_v(e) \cdot \partial_v(f),$$

where π is any uniformizer of the valuation ring $\mathcal{O}_v$ and $\{-1\}$ refers to the left-K_1^M-module structure of axiom **D3**. This verifies the well-definedness. The morphism of sheaves $\mathcal{M}_* \otimes_{\underline{\mathbf{Z}}} \mathcal{M}_* \to \mathcal{M}_{*+*}$ induces functorially a morphism between Čech cochain groups and composing this with the cup product, we get morphisms

$$\check{\mathrm{C}}^p(\mathfrak{U}, \mathcal{M}_n) \otimes_{\mathbf{Z}} \check{\mathrm{C}}^q(\mathfrak{U}, \mathcal{M}_m) \overset{\smile}{\longrightarrow} \check{\mathrm{C}}^{p+q}(\mathfrak{U}, \mathcal{M}_n \otimes_{\underline{\mathbf{Z}}} \mathcal{M}_m) \overset{\cdot}{\longrightarrow} \check{\mathrm{C}}^{p+q}(\mathfrak{U}, \mathcal{M}_{n+m}),$$

depending on the chosen pairing of cycle modules. This induces a morphism of Čech cohomology groups

$$\check{H}^p(\mathfrak{U}, \mathcal{M}_n) \otimes_{\mathbf{Z}} \check{H}^q(\mathfrak{U}, \mathcal{M}_m) \longrightarrow \check{H}^{p+q}(\mathfrak{U}, \mathcal{M}_{n+m}).$$

7.2. Cup Product on Idèles.

Next, we shall give a similar construction defining a pairing $\mathbf{I}_1 \otimes_{\mathbf{Z}} \mathbf{I}_1 \to \mathbf{I}_2$, which induces a pairing

$$H^1(\Gamma(X, \mathbf{I}_\bullet)) \otimes_{\mathbf{Z}} H^1(\Gamma(X, \mathbf{I}_\bullet)) \longrightarrow H^2(\Gamma(X, \mathbf{I}_\bullet)).$$

---

[8] Then the formula in eq. 7.2 need not hold for indices which are not strictly increasingly ordered, e.g. for $f \in \check{\mathrm{C}}^0$ and $g \in \check{\mathrm{C}}^1$, $I = \{1, 2\}$ one has $f_1 \cdot g_{12} = (f \smile g)_{12} = -(f \smile g)_{21}$, but this would $= -f_2 \cdot g_{21} = f_2 \cdot g_{12}$, which need not be true.

Explicitly, define a $\underline{\mathbf{Z}}$-bilinear pairing of sheaves of abelian groups

$$\vee : \mathbf{I}_1 \otimes \mathbf{I}_1 \to \mathbf{I}_2$$

$$e \otimes f \mapsto e \vee f \qquad \text{with } (e \vee f)_{x,y_i'} := (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} e^{01}_y) \cdot f^{12}_{x,y_i'}, \tag{7.3}$$

$$\text{(where } y \text{ is the image of } y_i' \text{ under } \operatorname{Spec} \widehat{\mathcal{O}}_x \to \operatorname{Spec} \mathcal{O}_x\text{)}$$

where again a cycle module is fixed and '$\cdot$' refers to a pairing as in eq. 7.1. Similarly to the cup product in eq. 7.2 this pairing does not have much symmetry before going to cohomology.

We now claim that on the level of cohomology, the above definition is the precise counterpart of the cup product construction on Čech cochains:

Proposition 11 (Product Compatibility). *Suppose X is integral and $\mathcal{M}_*$ admits a product structure (in the sense of eq. 7.1). Then the map $\vee$ descends to cohomology, i.e. is well-defined as in the lower row of*

(7.4)

$$\begin{tikzcd}
H^1(X, \mathcal{M}_*) \arrow[d] & \otimes_{\mathbf{Z}} & H^1(X, \mathcal{M}_*) \arrow[r, "\smile"] \arrow[d] & H^2(X, \mathcal{M}_*) \arrow[r, "\lambda"] & A^2(X, M_*) \arrow[d, equal] \\
H^1(\Gamma(X, \mathbf{I}_\bullet)) & \otimes_{\mathbf{Z}} & H^1(\Gamma(X, \mathbf{I}_\bullet)) \arrow[r, "\vee"'] & H^2(\Gamma(X, \mathbf{I}_\bullet)) \arrow[r, "\theta"'] & A^2(X, M_*)
\end{tikzcd}$$

and this diagram commutes, where the downward arrows and λ, θ are the comparison maps of Prop. 8 and Prop. 10 respectively.

Proof. **(1)** Firstly, we prove that the pairing in the lower row is well-defined, i.e. $\delta e \vee f$ and $e \vee \delta f$ are mapped to coboundaries. We only prove this for $e \vee \delta f$ and leave the other case to the reader: Let any two idèles

$$e = (e^{01}_y, e^{02}_x, e^{12}_{x,y_i'}) \in \Gamma(X, \mathbf{I}_1) \qquad \text{and} \qquad f = (f^0_\eta, f^1_y, f^2_x) \in \Gamma(X, \mathbf{I}_0)$$

be given. With the definition in eq. 7.3 we obtain

$$(\delta f \vee e)^{012}_{x,y_i'} := (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} f^1_y - \operatorname{res}^{\widehat{K}_{x,y_i'}}_{K_\eta} f^0_\eta) \cdot e^{12}_{x,y_i'}. \tag{7.5}$$

As usual, y denotes the image of y_i' under $\operatorname{Spec} \widehat{\mathcal{O}}_x \to \operatorname{Spec} \mathcal{O}_x$. Next, define $j = (j^{01}_y, j^{02}_x, j^{12}_{x,y_i'}) \in \Gamma(X, \mathbf{I}_1)$ by

$$j^{01}_y = (\operatorname{res}^{\widehat{K}_y}_{K_\eta} f^0_\eta) \cdot e^{01}_y \qquad j^{02}_x = (\operatorname{res}^{\widehat{K}_x}_{K_\eta} f^0_\eta) \cdot e^{02}_x \qquad j^{12}_{x,y_i'} = (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} f^1_y) \cdot e^{12}_{x,y_i'}.$$

Then the idelic differential yields (by using axiom **P2a** of a cycle module pairing several times, eq. 7.5 and $\delta e = 0$ which we may assume since we only need our claim for cocycles)

$$\begin{aligned}
(\delta j)^{012}_{x,y_i'} &= j^{12}_{x,y_i'} - \operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_x} j^{02}_x + \operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} j^{01}_y \\
&= (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} f^1_y) \cdot e^{12}_{x,y_i'} - (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{K_\eta} f^0_\eta) \cdot (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_x} e^{02}_x - \operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} e^{01}_y) \\
&= (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{\widehat{K}_y} f^1_y) \cdot e^{12}_{x,y_i'} - (\operatorname{res}^{\widehat{K}_{x,y_i'}}_{K_\eta} f^0_\eta) \cdot e^{12}_{x,y_i'} = (\delta f \vee e)^{012}_{x,y_i'}
\end{aligned}$$

In particular, $\delta f \vee e$ is a coboundary. For $e \vee \delta f$ one proceeds similarly.

(2) Next, we prove the commutativity of eq. 7.4. Suppose $(g_{\alpha\beta})_{\alpha\beta}, (h_{\alpha\beta})_{\alpha\beta} \in \check{C}^1(\mathfrak{U}, \mathcal{M}_*)$ are Čech 1-cocycle representatives of arbitrary elements $g, h \in H^1(X, \mathcal{M}_*)$

and a sufficiently fine open cover $\mathfrak{U}$. According to the description of the cup product in eq. 7.2, the image in $\check{H}^2(\mathfrak{U}, \mathcal{M}_*)$ in the upper row is represented by the Čech 2-cocycle

$$(g \smile h)_{\alpha\beta\gamma} = {}^{\mathcal{M}_*}\operatorname{res}^{U_{\alpha\beta}}_{U_{\alpha\beta\gamma}} g_{\alpha\beta} \cdot {}^{\mathcal{M}_*}\operatorname{res}^{U_{\beta\gamma}}_{U_{\alpha\beta\gamma}} h_{\beta\gamma} \in \check{\mathrm{C}}^2(\mathfrak{U}, \mathcal{M}_*),$$

where ${}^{\mathcal{M}_*}\operatorname{res}^{U_{\alpha\beta}}_{U_{\alpha\beta\gamma}}$ denotes the restriction map of the sheaf $\mathcal{M}_*$ and is not to be confused with the restrictions of the cycle module M_*. We fix a disjoint decomposition $X = \bigcup_{\alpha\in I} \Sigma_\alpha$ as in Lemma 17. According to Prop. 9 idèle representatives for the image under $\tau : \check{H}^1(\mathfrak{U}, \mathcal{M}_*) \to H^1(\Gamma(X, \mathbf{I}_\bullet))$, $g' := \tau g$, $h' := \tau h$, are given by Čech 0-cocycles in

$$g'^{01}_y := -\operatorname{res}^{\widehat{K}_y}_{K_\eta} g_{\beta_\eta\beta_y} \qquad g'^{02}_x := -\operatorname{res}^{\widehat{K}_x}_{K_\eta} g_{\beta_\eta\beta_x} \qquad g'^{12}_{x,y'_i} := -\operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} g_{\beta_y\beta_x},$$

using the same notation as in the statement of Prop. 9, analogously for h'. Using these representatives and writing '$\cdot$' for the pairing of the cycle modules, the image in the lower row under $\vee$ (see eq. 7.3) is

$$\begin{aligned}(g' \vee h')_{x,y'_i} &= (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} g'^{01}_y) \cdot h'^{12}_{x,y'_i} \in H^2(X, \mathbf{I}_\bullet) \\ &= (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} g_{\beta_\eta\beta_y}) \cdot (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} h_{\beta_y\beta_x}) = \operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} (g_{\beta_\eta\beta_y} \cdot h_{\beta_y\beta_x})\end{aligned}$$

by **R1a**, **P2a**. According to Prop. 8 the image in $A^2(X, M_*)$ can be represented by

$$\begin{aligned}\rho_x &= \left(\sum\nolimits_{y'_i} \partial^{y'_i}_x \partial_{y'_i} (g' \vee h')_{x,y'_i}\right)_x \in \coprod\nolimits_{x\in X^{(2)}} M_{*-2}(\kappa(x)) \\ &= \sum\nolimits_{y'_i} \partial^{y'_i}_x \partial_{y'_i} (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{K_\eta} (g_{\beta_\eta\beta_y} \cdot h_{\beta_y\beta_x})). \qquad (7.6)\end{aligned}$$

For every closed point $x \in X^{(2)}$ the set $\{y'_1, \dots, y'_r\}$ gives the fibers of a height 1 prime y of $\mathcal{O}_x$ under $\operatorname{Spec} \widehat{\mathcal{O}}_x \to \operatorname{Spec} \mathcal{O}_x$. For the upper row in eq. 7.4 we use Prop. 10 and obtain

$$\begin{aligned}\rho'_x &= \sum_y \partial^y_x \partial^\eta_y (g \smile h)_{\beta_\eta\beta_y\beta_x} \\ &= \sum_y \partial^y_x \partial^\eta_y ({}^{\mathcal{M}_*}\operatorname{res}^{U_{\beta_\eta\beta_y}}_{U_{\beta_\eta\beta_y\beta_x}} g_{\beta_\eta\beta_y} \cdot {}^{\mathcal{M}_*}\operatorname{res}^{U_{\beta_y\beta_x}}_{U_{\beta_\eta\beta_y\beta_x}} h_{\beta_y\beta_x}).\end{aligned}$$

For an open U the group $\mathcal{M}_*(U)$ is a subgroup of $M_*(\kappa(\eta))$, so we may consider $g_{\beta_\eta\beta_y}, h_{\beta_y\beta_x}$ as elements of $M_*(\kappa(\eta))$, η being the *unique* generic point of X, and with this interpretation we can discard the restrictions à la ${}^{\mathcal{M}_*}\operatorname{res}^{U_{\beta_\eta\beta_y}}_{U_{\beta_\eta\beta_y\beta_x}}$ in the above. Comparing this equality with the one in eq. 7.6, our claim follows from using the commutativity of the diagram

$$\begin{tikzcd}
M_*(\widehat{K}_{x,y'_i}) \arrow[r, "\partial_{y'_i}"] & M_{*-1}(\kappa(y'_i)) \arrow[d, shift left] & \\
M_*(\widehat{K}_x) \arrow[u] \arrow[r, "\partial_{y'}"] & \coprod_{y'\in(\widehat{\mathcal{O}}_x)^{(1)}} M_{*-1}(\kappa(y')) \arrow[u, shift left] \arrow[r, "\partial_x"] & M_{*-2}(\kappa(x)) \arrow[d, equal] \\
M_*(K_\eta) \arrow[u] \arrow[r, "\partial_y"'] & \coprod_{y\in(\mathcal{O}_x)^{(1)}} M_{*-1}(\kappa(y)) \arrow[u] \arrow[r, "\partial_x"'] & M_{*-2}(\kappa(x)),
\end{tikzcd}$$

where the upward arrows are the flat pullbacks induced from $\operatorname{Spec}\widehat{\mathcal{O}}_x \to \operatorname{Spec}\mathcal{O}_x$ as well as $\operatorname{Spec}\widehat{\mathcal{O}}_{x,y_i'} \to \operatorname{Spec}\widehat{\mathcal{O}}_x$ (these morphisms are flat of constant relative dimension by Lemma 2), and the single downward arrow is the canonical splitting of the upward morphism. This finishes the proof. $\square$

The previous result immediately yields a formula for intersection multiplicities on surfaces. We use that the product of the Chow ring of a smooth proper surface over a field k coincides with the cup product pairing of the corresponding sheaf cohomology groups $H^p(X, \mathcal{K}_p^M)$:

$$\begin{tikzcd}
\mathrm{CH}^p(X) \arrow[d] & \otimes_{\mathbf{Z}} & \mathrm{CH}^q(X) \arrow[d] \arrow[r, "\alpha\cdot\beta"] & \mathrm{CH}^{p+q}(X) \arrow[d] \\
H^p(X, \mathcal{K}_p^M) & \otimes_{\mathbf{Z}} & H^q(X, \mathcal{K}_q^M) \arrow[r, "{[\alpha]\smile[\beta]}"'] & H^{p+q}(X, \mathcal{K}_{p+q}^M).
\end{tikzcd} \tag{7.7}$$

For algebraic K-theory sheaves $\mathcal{K}_n$ this was proven by Grayson, following Bloch and Quillen, see [**13**, Thm. 1.2] (the general version for cycle modules is by Rost and in [**26**]); depending on conventions there is a sign twist $(-1)^{pq}$, see [**13**].

Proposition 12 (Intersection Multiplicity Formula, [**24**] Cor. to Prop. 2)**.** *Let X be an integral surface which is smooth proper over a field k. Let Z_1, Z_2 be two irreducible divisors. For each Z_i, $i = 1, 2$, let*

(1) $f_y^{(i)} := \begin{cases} \text{any uniformizer of } \widehat{\mathcal{O}}_y & \text{for } y = y_{Z_i} \\ \text{any unit in } \widehat{\mathcal{O}}_y^{\times} & \text{for } y \neq y_{Z_i}\text{; and} \end{cases}$

(2) $g_x^{(i)} := \operatorname{res}_{K_\eta}^{\widehat{K}_x} \begin{cases} \text{any principal ideal gen. of } y_{Z_i} \text{ in } \mathcal{O}_x & \text{for } x \in \overline{\{y_{Z_i}\}} \\ \text{any unit in } \mathcal{O}_x^{\times} & \text{for } x \notin \overline{\{y_{Z_i}\}}. \end{cases}$

Then the intersection multiplicity is given by

$$Z_1 \cdot Z_2 = \sum\nolimits_{x,y_i'} [\kappa(x) : k] \cdot \partial_x^{y_i'} \partial_{y_i'} \{\operatorname{res}_{\widehat{K}_y}^{\widehat{K}_{x,y_i'}} f_y^{(1)}, \operatorname{res}_{\widehat{K}_x}^{\widehat{K}_{x,y_i'}} g_x^{(2)}\}, \tag{7.8}$$

where the sum runs over all closed points $x \in X^{(2)}$ and y_i' over all non-transcendental primes of $\widehat{\mathcal{O}}_x$; y is the image of y_i' under $\operatorname{Spec}\widehat{\mathcal{O}}_x \to \operatorname{Spec}\mathcal{O}_x$. The sum has only finitely many non-zero terms.

Remark 6. *It is noteworthy that while the symbol $\{x, y\}$ is anti-commutative in its slots, the intersection pairing $Z_1 \cdot Z_2$ (and generally the product in the Chow ring) is symmetric.*

Proof. The $\mathbf{Z}$-valued intersection pairing $Z_1 \cdot Z_2$ is defined as the composition

$$\mathrm{CH}^1(X) \otimes \mathrm{CH}^1(X) \xrightarrow{\cdot} \mathrm{CH}^2(X) \xrightarrow{j_*} \mathrm{CH}_0(\operatorname{Spec} k) \cong \mathbf{Z},$$

where $j : X \to \operatorname{Spec} k$ is the structure morphism. Using the commutativity of eq. 7.7 and combining it with Prop. 11, we may equivalently evaluate

$$H^1(\Gamma(X, \mathbf{I}_\bullet)) \otimes H^1(\Gamma(X, \mathbf{I}_\bullet)) \longrightarrow H^2(\Gamma(X, \mathbf{I}_\bullet)).$$

Firstly, we use Prop. 7 for the (Milnor) K-cycle module $M_* = K_*^M$, so in degree 1 we get an explicit description of the map $\mathrm{CH}^1(X) =_{\text{def.}} A^1(X, K_1^M) \to H^1(\Gamma(X, \mathbf{I}_\bullet))$, specifically for Z_1 we get $\alpha := (\alpha_y^{01}, \alpha_x^{02}, \alpha_{x,y_i'}^{12}) \in \mathbf{I}_1$:

$$\alpha_y^{01} = f_y, \qquad \alpha_x^{02} = g_x, \qquad \alpha_{x,y_i'}^{12} = \operatorname{res}_{\widehat{K}_x}^{\widehat{K}_{x,y_i'}} g_x - \operatorname{res}_{\widehat{K}_y}^{\widehat{K}_{x,y_i'}} f_y.$$

Note that the explicit construction of g_x is particularly simple in this case since $K^M_{-1}(F) = 0$, compare with eq. 5.2. Analogously for for Z_2, $\beta := (\beta^{01}_y, \beta^{02}_x, \beta^{12}_{x,y'_i})$. Then we get $(\alpha \vee \beta)_{x,y'_i} \in H^2(\Gamma(X, \mathbf{I}_\bullet))$ and using Prop. 8 this maps to $\mathrm{CH}^2(X) =_{\text{def.}} A^2(X, K^M_2)$ as

$$\left(\sum\nolimits_{y'_i} \partial^{y'_i}_x \partial_{y'_i} (\alpha \vee \beta)_{x,y'_i} \right)_x \in \coprod\nolimits_{x \in X^{(2)}} M_{*-2}(\kappa(x))$$

and unwinding the proper push-forward j_* yields

$$Z_1 \cdot Z_2 = \sum\nolimits_{x,y'_i} \operatorname{cor}^{\kappa(x)}_k \partial^{y'_i}_x \partial_{y'_i} \{ \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} \alpha^{01}_y, \beta^{12}_{x,y'_i} \},$$

and last bracket unwinds as $\{ \operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} f_y, (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_x} g_x) \cdot (\operatorname{res}^{\widehat{K}_{x,y'_i}}_{\widehat{K}_y} f_y)^{-1} \}$. We can discard the term of the shape $\{f_y, f_y^{-1}\}$ since it is 2-torsion and thus when sent to $\mathbf{Z}$ must go to zero necessarily. For K^M_0 the corestriction is just multiplication with the extension degree, giving eq. 7.8. □

Remark 7. *A straight-forward variation of the above proof yields the formula eq. 0.2 by replacing the cup product of the idèle resolution by the cup product of the Čech resolution.*

8. Example with negative self-intersection

Finally, we wish to give an example of an explicit computation of a negative self-intersection number using the methods of this text. No moving will be used. To keep the example sufficiently simple, we consider Hirzebruch surfaces F_n ($n \in \mathbf{Z}$). They can be described either as the toric surface of the fan (see [**6**] §1-3 for toric generalities)

$$\sigma_0 = \mathbf{R}e_x + \mathbf{R}e_y \qquad \sigma_1 = \mathbf{R}e_y + \mathbf{R}(-e_x + ne_y)$$
$$\sigma_2 = \mathbf{R}(-e_x + ne_y) - \mathbf{R}e_y \qquad \sigma_3 = \mathbf{R}e_x - \mathbf{R}e_y$$





(depicted left for $n = 2$) or as the projectivization of the vector bundle $\pi : E \to \mathbf{P}^1_k$ given through $\mathcal{O}_{\mathbf{P}^1_k}(n) \oplus \mathcal{O}_{\mathbf{P}^1_k}$ on $\mathbf{P}^1_k$. For $n = 0$ one has $F_0 \simeq \mathbf{P}^1_k \times \mathbf{P}^1_k$. Explicitly, the smooth proper surface F_n is patched from affine opens

$$U_0 = \operatorname{Spec} k[X, Y] \qquad U_1 = \operatorname{Spec} k[X^{-1}, X^n Y]$$
$$U_2 = \operatorname{Spec} k[X^{-1}, X^{-n} Y^{-1}] \qquad U_3 = \operatorname{Spec} k[X, Y^{-1}]$$

(all isomorphic to $\mathbf{A}^2_k$) along the intersections

$$U_{01} = \operatorname{Spec} k[X^{-1}, X, Y] \qquad U_{12} = \operatorname{Spec} k[X^{-n} Y^{-1}, X^n Y, X^{-1}]$$
$$U_{23} = \operatorname{Spec} k[X^{-1}, X, Y^{-1}] \qquad U_{03} = \operatorname{Spec} k[X, Y, Y^{-1}]$$

(all isomorphic to $\mathbf{A}^1_k \times \mathbf{G}_{m,k}$). All other intersections of two opens, e.g. U_{02} or U_{13}, are 2-tori, $\operatorname{Spec} k[X, X^{-1}, Y, Y^{-1}] \simeq \mathbf{G}^2_{m,k}$. The same holds for all triple intersections like U_{012}, U_{013}, etc. These opens U_i correspond to the toric affine opens coming from cones σ_i. In particular, $\mathfrak{U} := (U_i)_{i=0,1,2,3}$ is an open cover we may use for the Čech cohomology. Define $V := F_n \setminus U_0$ as the reduced closed complement of U_0. Using the same cover as for F_n, this locally comes down to

$$V_0 = U_0 \setminus U_0 = \varnothing, \quad V_1 = U_1 \setminus U_0 = \operatorname{Spec} k[X^nY], \quad V_3 = U_3 \setminus U_0 = \operatorname{Spec} k[X],$$
$$V_2 = U_2 \setminus D_{X^{-1}(X^{-n}Y^{-1})} = \operatorname{Spec} k[X^{-1}, X^{-n}Y^{-1}]/(X^{-1}) \cdot (X^{-n}Y^{-1});$$

in particular $V_1 \simeq \mathbf{A}^1_k$, $V_3 \simeq \mathbf{A}^1_k$, V_2 is reducible and its irreducible components are both isomorphic to $\mathbf{A}^1_k$. The closures (in the whole surface F_n) $\overline{V_1}$ and $\overline{V_3}$ are both isomorphic to $\mathbf{P}^1_k$ (with $V_{12} \simeq \mathbf{G}_{m,k}$ and $V_{23} \simeq \mathbf{G}_{m,k}$ the overlaps of the two copies of $\mathbf{A}^1_k$), they intersect in a single closed point in V_2. Moreover, $V_{13} = \varnothing$. For the construction of the contracting homotopy in Lemma 17 we now may use the disjoint decomposition of F_n (as a set!)

$$\Sigma_0 := U_0, \quad \Sigma_1 := V \cap U_1, \quad \Sigma_2 := (V \setminus U_1) \cap U_2, \quad \Sigma_3 := (V \setminus (U_1 \cup U_2)).$$

Graphically, the decomposition of the complement V is depicted on the right in the above figure. The two circles represent $\overline{V_1}$ and $\overline{V_3}$ ($\simeq \mathbf{P}^1_k$). Summarized, F_n decomposes as follows:

- (height 2) the unique generic point η lies in Σ_0;
- (height 1) the generic points of all integral curves of U_0 are in Σ_0. $\Sigma_1^{(1)}$ contains only the height 1 generic point of $\overline{V_1}$, $\Sigma_2^{(1)}$ contains only the height 1 generic point of $\overline{V_3}$, Σ_3 does not contain any height 1 points;
- (height 0) the closed points of U_0 all lie in Σ_0. The closed points in Σ_1 are the closed points of $V_1 \simeq \mathbf{A}^1_k$ (the single additional closed point of its closure $\overline{V_1}$ lies in Σ_2 - it's the same point as the intersection $\overline{V_1} \cap \overline{V_3}$), the closed points in Σ_2 are the closed points of $V_3 \simeq \mathbf{A}^1_k$ (the single additional closed point of its closure $\overline{V_3}$ lies in Σ_3). Σ_3 consists only of this closed point.

In particular, the set-theoretic intersection closed point of $\overline{V_1}$ and $\overline{V_3}$ lies in Σ_2.

Now we wish to study the divisor D associated to the cone spanned by e_y, i.e. $\sigma_0 \cap \sigma_1$.

Claim: We have self-intersection $D \cdot D = -n$.

The divisor D is an effective Cartier divisor in $H^0(F_n, \mathcal{K}^\times/\mathcal{O}^\times)$, which we may represent in our Čech cover through

$$c_0 = Y \qquad c_1 = X^nY \qquad c_2 = 1 \qquad c_3 = 1$$

in $\check{H}^0(\mathfrak{U}, \mathcal{K}^\times/\mathcal{O}^\times)$. Our cover is fine enough to admit the morphism $\check{H}^0(\mathfrak{U}, \mathcal{K}^\times/\mathcal{O}^\times) \to \check{H}^1(\mathfrak{U}, \mathcal{O}^\times)$, giving the associated line bundle

$$\tilde{c}_{01} = X^n \qquad \tilde{c}_{02} = Y^{-1} \qquad \tilde{c}_{03} = Y^{-1}$$
$$\tilde{c}_{12} = X^{-n}Y^{-1} \qquad \tilde{c}_{13} = X^{-n}Y^{-1} \qquad \tilde{c}_{23} = 1.$$

Thanks to Prop. 10 and the diagram of eq. 7.7 the self-intersection number $D \cdot D$ comes down to the computation of various tame symbols:

$$D \cdot D = j_* h \qquad \text{with } h_x = j_* \sum_{x \in \overline{\{y\}}} \partial^y_x \partial^\eta_y (\tilde{c} \smile \tilde{c})_{\beta_\eta \beta_y \beta_x} \in \coprod_{x \in X^{(2)}} \mathbf{Z}$$
$$= j_* \sum_{x,y} \partial^y_x \partial^\eta_y \{\tilde{c}_{\beta_\eta \beta_y}, \tilde{c}_{\beta_y \beta_x}\} \tag{8.1}$$

($j : X \to \operatorname{Spec} k$ is the structure morphism; compare with eq. 7.8; the meaning of $\beta_\eta, \beta_y, \beta_x$ is as in Prop. 10; see also eq. 0.2). As the formula eq. 6.2 really mostly depends on the values of $\beta_\eta, \beta_y, \beta_x$ for various y, x, it is convenient to do a case-distinction depending on these values. Since $\beta_\eta = 0$ always, we are left with

$$\{\tilde{c}_{\beta_\eta\beta_y}, \tilde{c}_{\beta_y\beta_x}\} = \left\{ \begin{array}{|l||l|l|l|l|} \hline \beta_x \ / \ \beta_y & 0 & 1 & 2 & 3 \\ \hline\hline 0 & 0 & 0 & * & * \\ \hline 1 & 0 & 0 & \{Y^{-1}, X^n Y\} & \{Y^{-1}, X^n Y\} \\ \hline 2 & 0 & \{X^n, X^{-n}Y^{-1}\} & 0 & 0 \\ \hline 3 & 0 & \{X^n, X^{-n}Y^{-1}\} & 0 & 0, \\ \hline \end{array} \right.$$

where $*$ indicates an element of the shape $\{a, a^{-1}\}$. Whereas these elements are usually non-zero, we have $\{a, a^{-1}\} = \{a, -1\}$, so they are 2-torsion. Thus, when being mapped to an intersection number, i.e. to $\mathbf{Z}$, they necessarily vanish, so we may disregard them already here. The value $\beta_y = 3$ is impossible since Σ_3 does not contain generic points of curves. The value $\beta_y = 2$ is only possible if $y = \overline{V_3}$, but the only non-trivial entry is at $\beta_x = 1$, however by the nature of our decomposition no closed points on the curve $\overline{V_3}$ lie in Σ_1. Thus, only for $\beta_y = 1$ non-trivial symbols occur. Note that $\beta_y = 1$ implies $y = \overline{V_1}$ and all the closed points of $\overline{V_1}$ lie in Σ_1 and Σ_2, so the case $\beta_x = 3$ is also impossible. For $\beta_x = 2$ the closed point must be $\overline{V_1} \cap \overline{V_3}$, given by $x = (X^{-1}, X^{-n}Y^{-1})$ in $U_2 = \operatorname{Spec} k[X^{-1}, X^{-n}Y^{-1}]$ and $y \mid_{U_2} = \overline{V_1} \mid_{U_2} = (X^{-1})$. Hence, the whole sum of eq. 8.1 reduces to the single expression

$$\begin{aligned} D \cdot D &= j_* \partial_x \partial_y \{X^n, X^{-n}Y^{-1}\} = j_* \partial_x \partial_y (-n\{X^{-1}, X^{-n}Y^{-1}\}) \\ &= -n j_* \partial_x \{\overline{X^{-n}Y^{-1}}\} = -n j_* [x, 1_{\mathbf{Z}}] = -n \in \mathbf{Z} \end{aligned}$$

since x is a closed point of degree 1 on F_n.

9. Perspectives

Apart from cycle modules various other 'coefficient systems' for adèles/idèles can be considered, e.g. the aforementioned case of quasi-coherent sheaves or Gorchinskiy's homology sheaves [**12**]. Apart from encoding the respective cohomology groups, these constructions sometimes also allow very interesting approaches to prove various results, e.g. the residue theorem on curves as pioneered by Tate [**29**] or Weil's Reciprocity Law for tame symbols on curves, see [**2**]. A related approach can for example be found in [**1**].

9.1. The Anticipated Picture for Schemes over Z. The strongest driving force behind the concept of adèles/idèles is however the hope to use these mechanisms in an arithmetic context, i.e. for schemes over $\operatorname{Spec} \mathbf{Z}$. We do not really touch this matter in the present text, but certainly this text concerns the first step in the following general philosophy:

- Formulate invariants of objects in a geometric context (over a field, or in positive characteristic where no archimedean places are to be expected) using the language of idèles.
- Then transfer such an idèle picture to an arithmetic situation, e.g. a number field or an arithmetic surface $X \to \operatorname{Spec} \mathbf{Z}$ and exploit that the language of idèles seems to be predestined for implementing archimedean contribution.

At least for number fields Chevalley's idèles (including infinite places) provide a remarkably powerful tool. It is reasonable to expect that a similar approach for arithmetic surfaces can also be carried out. The key point is that it seems to be difficult to implement a theory of archimedean places using Zariski sheaves on schemes since the archimedean points are not represented by points of the space underlying a scheme. One hopes that direct manipulations with idèle complexes can provide a more natural setup for archimedean investigations. We quickly want to outline a sketch of the expected picture for an integral proper flat surface $j : X \rightarrow \operatorname{Spec} \mathbf{Z}$. The set of codimension 1 points y decomposes disjointly into *vertical* and *horizontal* curves, according to whether the image $j(y)$ is of codimension 1 or 0. Then the horizontal curves $i : \overline{\{y\}} \hookrightarrow X$ are orders in number fields in a canonical way by considering the composition $j \circ i : \overline{\{y\}} \rightarrow \operatorname{Spec} \mathbf{Z}$. Hence, there is a natural notion of an infinite place for horizontal curves which should play the role of additional closed points.

9.2. Relations to L-Functions. One of the most intriguing questions (raised by Parshin, [**22**]) is whether it is possible to extend the Tate-Iwasawa method to 2-dimensional adèles & idèles in order to treat zeta and L-functions of arithmetic surfaces on a similar footing as in the case of number fields. See the work of Fesenko [**10**], [**11**] for this matter. Various problems need to be (and have been) overcome:
(A) The involved local objects, e.g. $\widehat{\mathcal{O}}_x^\times$, need to be equipped with a topology. For $\widehat{\mathcal{O}}_x^\times$ this is straight-forward, but for groups like $K_2(\widehat{K}_x)$ or $K_2(\widehat{K}_{x,y})$ the matter is considerably more complicated. However, answers to these problems are known and we survey this in §2.3. For a general cycle module M_* and a topological field F the question of equipping the groups $M_n(F)$ with a suitable topology is completely open.
(B) In order to have harmonic analysis available, these groups would classically have to be locally compact to carry a Haar measure. This is fine for classical 1-dimensional idèles (by the assumption of finite residue fields), but for surfaces the corresponding groups are *not* locally compact; no Haar measure exists. We again refer to A. Weil's [**31**, p. VI (Foreword), 2nd paragraph] for a discussion strongly suggesting that this is rather a defect of an insufficiently general harmonic analysis than an actual organic obstacle. For the construction of 2-dimensional zeta integrals a generalized harmonic analysis needs to be used. See [**11**] for details on this approach using K^t-idèles (the groups K^t reappear in §2.3 in this text). See also [**18**] for 2-dimensional Haar integration, or [**14**] for more on this circle of problems. Many open questions remain though.
(C) Following this line of thought, one is led to the observation that instead of the (so called rank 1) ring of integers $\widehat{\mathcal{O}}_{x,y} \subset \widehat{K}_{x,y}$ in the 2-dimensional local fields $\widehat{K}_{x,y}$ is probably not the correct object to consider for harmonic analysis purposes. Rather, the (so called rank 2) ring of integers $O_{x,y}$ turns out to be the relevant object. This ring arises as the (Krull) discrete valuation ring on the right-hand side in

$$\begin{array}{cc} \widehat{\mathcal{O}}_{x,y}^\times \rightarrow \widehat{K}_{x,y}^\times \overset{v_1}{\rightarrow} \mathbf{Z} & \qquad O_{x,y} \rightarrow \widehat{K}_{x,y}^\times \overset{v_2}{\rightarrow} \mathbf{Z} \oplus \mathbf{Z} \\ \text{value group } \mathbf{Z} & \qquad \text{value group } \mathbf{Z}^{\oplus 2}, \end{array}$$

where v_2 is a lexicographically ordered rank 2 discrete valuation coming from both the curve y and the point x. In the present text we have not touched this perspective. See [**11**] for more information.

References

1. Greg W. Anderson and Fernando Pablos Romo, *Simple proofs of classical explicit reciprocity laws on curves using determinant groupoids over an Artinian local ring.*, Commun. Algebra **32** (2004), no. 1, 79–102 (English).
2. E. Arbarello, C. De Concini, and V.G. Kac, *The infinite wedge representation and the reciprocity law for algebraic curves.*, Theta functions, Proc. 35th Summer Res. Inst. Bowdoin Coll., Brunswick/ME 1987, Proc. Symp. Pure Math. 49, Pt. 1, 171-190 (1989)., 1989.
3. A.A. Beĭ linson, *Residues and adeles.*, Funct. Anal. Appl. **14** (1980), 34–35 (English. Russian original).
4. Spencer Bloch, *K_2 and algebraic cycles*, Ann. of Math. (2) **99** (1974), 349–379. MR 49 #7260
5. Spencer Bloch and Arthur Ogus, *Gersten's conjecture and the homology of schemes*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 181–201 (1975). MR 54 #318
6. V. I. Danilov, *The geometry of toric varieties*, Russian Mathematical Surveys (1978), 33(2):97.
7. Frédéric Déglise, *Transfers on Chow groups with coefficients. (Transferts sur les groupes de Chow à coefficients.)*, Math. Z. **252** (2006), no. 2, 315–343 (French).
8. Elman, R. and Karpenko, N. and Merkurjev, A., *The algebraic and geometric theory of quadratic forms*, American Mathematical Socienty Colloquium Publications (Vol 56), AMS, Providence, RI, 2008.
9. Ivan Fesenko, *Sequential topologies and quotients of Milnor K-groups of multidimensional local fields.*, St. Petersbg. Math. J. **13** (2002), no. 3, 485–501 (English. Russian original).
10. ———, *Adelic approach to the zeta function of arithmetic schemes in dimension two.*, Mosc. Math. J. **8** (2008), no. 2, 273–317 (English).
11. ———, *Analysis on arithmetic schemes. II.*, (2010) (English).
12. S.O. Gorchinskii, *An adelic resolution for homology sheaves.*, Izv. Math. **72** (2008), no. 6, 1187–1252 (English. Russian original).
13. Daniel R. Grayson, *Products in K-theory and intersecting algebraic cycles*, Invent. Math. **47** (1978), no. 1, 71–83. MR 58 #10890
14. Ehud Hrushovski and David Kazhdan, *Integration in valued fields.*, Ginzburg, Victor (ed.), Algebraic geometry and number theory. In Honor of Vladimir Drinfeld's 50th birthday. Basel: Birkhäuser. Progress in Mathematics 253, 261-405 (2006)., 2006.
15. A. Huber, *On the Parshin-Beilinson adeles for schemes.*, Abh. Math. Semin. Univ. Hamb. **61** (1991), 249–273 (English).
16. O. Izhboldin, *p-primary part of the Milnor K-groups and Galois cohomology of fields of characteristic p.*, Fesenko, Ivan (ed.) et al., Invitation to higher local fields. Extended version of talks given at the conference on higher local fields, Münster, Germany, August 29-September 5, 1999. Coventry: Geometry and Topology Publications, Geom. Topol. Monogr. 3, 19-29 (2000)., 2000.
17. M. Kurihara, *Kato's higher local class field theory.*, Fesenko, Ivan (ed.) et al., Invitation to higher local fields. Extended version of talks given at the conference on higher local fields, Münster, Germany, August 29-September 5, 1999. Coventry: Geometry and Topology Publications, Geom. Topol. Monogr. 3, 53-60 (2000)., 2000.
18. Matthew Morrow, *Integration on valuation fields over local fields.*, Tokyo J. Math. **33** (2010), no. 1, 235–281 (English).
19. Denis V. Osipov, *n-dimensional local fields and adeles on n-dimensional schemes.*, Young, Nicholas (ed.) et al., Surveys in contemporary mathematics. Cambridge: Cambridge University Press. London Mathematical Society Lecture Note Series 347, 131-164 (2008)., 2008.
20. D.V. Osipov, *Adele constructions of direct images of differentials and symbols.*, Sb. Math. **188** (1997), no. 5, 697–723 (English. Russian original).
21. Osipov, D. and Zhu, X., *A categorical proof of the Parshin reciprocity laws on algebraic surfaces*, arXiv:1002.4848, to appear in Algebra and Number Theory.
22. A. N. Parshin, *Representations of higher adelic groups and arithmetic*, arXiv:1012.0486v1 [math.NT].
23. A.N. Parshin, *On the arithmetic of two-dimensional schemes. I: Distributions and residues.*, (1977) (English).
24. ———, *Chern classes, adeles and L-functions.*, J. Reine Angew. Math. **341** (1983), 174–192 (English).

25. Daniel Quillen, *Higher algebraic K-theory. I*, Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Springer, Berlin, 1973, pp. 85–147. Lecture Notes in Math., Vol. 341. MR 49 #2895
26. Markus Rost, *Chow groups with coefficients*, Doc. Math. **1** (1996), No. 16, 319–393 (electronic). MR 98a:14006
27. J. P. Serre, *Algebraic groups and class fields*, GTM, Springer Verlag, New York, 1988.
28. Tate, J., *Fourier analysis in number fields and Heckes zeta-functions (doctoral thesis, 1950)*, Algebraic Number Theory (ed. by. J.W.S. Cassels, A. Frhlich), Academic Press, London, 1967.
29. ———, *Residues of differentials on curves.*, Ann. Sci. c. Norm. Supr. (4) 1, No. 1, 149-159 (1968). ISSN 0012-9593, 1968.
30. Wilberd van der Kallen, *The K_2 of rings with many units*, Ann. Sci. École Norm. Sup. (4) **10** (1977), no. 4, 473–515. MR 58 #22018
31. A. Weil, *Basic number theory*, Classics in Mathematics (reprint of the 1974 edition), Springer Verlag, New York, 1995.
32. I. Zhukov, *Milnor and topological K-groups of higher-dimensional complete fields*, Algebra i Analiz **9** (1997), no. 1, 98–147. MR 98m:11131

School of Mathematical Sciences, University of Nottingham, University Park, Nottingham, NG7 2RD, UK (Europe)

E-mail address: `pmxob2@nottingham.ac.uk`